

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 5/10, 15/00, 15/11	A1	(11) International Publication Number: WO 95/02686 (43) International Publication Date: 26 January 1995 (26.01.95)
(21) International Application Number: PCT/US94/06675 (22) International Filing Date: 14 June 1994 (14.06.94) (30) Priority Data: 08/093,210 16 July 1993 (16.07.93) US (71) Applicant: THE GENERAL HOSPITAL CORPORATION [US/US]; 55 Fruit Street, Boston, MA 02114 (US). (72) Inventors: SEED, Brian; Nine Hawthorne Place, #5J, Boston, MA 02114 (US). ROMEO, Charles; 45 Chester Road, Belmont, MA 02178 (US). KOLANUS, Waldemar; 81 Templeton Parkway, Watertown, MA 02172 (US). (74) Agent: CLARK, Paul, T.; Fish & Richardson, 225 Franklin Street, Boston, MA 02110-2804 (US).		(81) Designated States: AU, CA, CZ, FI, HU, JP, KR, NO, NZ, RU, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: REDIRECTION OF CELLULAR IMMUNITY BY RECEPTOR CHIMERAS (57) Abstract Disclosed is a method of directing a cellular response in a mammal by expressing in a cell of the mammal a chimeric receptor which causes the cells to specifically recognize and destroy an infective agent, a cell infected with an infective agent, a tumor or cancerous cell, or an autoimmune-generated cell. The chimeric receptor includes an extracellular portion which is capable of specifically recognizing and binding the target cell or target infective agent, and (b) an intracellular portion of a protein-tyrosine kinase which is capable of signalling the therapeutic cell to destroy a receptor-bound target cell or a receptor-bound target infective agent. Also disclosed are cells which express the chimeric receptors and DNA encoding the chimeric receptors.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

REDIRECTION OF CELLULAR IMMUNITY BY RECEPTOR CHIMERASField of the Invention

The invention concerns functional protein-tyrosine
5 kinase chimeras which are capable of redirecting immune
system function. More particularly, it concerns the
regulation of lymphocytes, macrophages, natural killer
cells or granulocytes by the expression in said cells of
chimeras which cause the cells to respond to targets
10 recognized by the chimeras. The invention also concerns
functional protein-tyrosine kinase chimeras which are
capable of directing therapeutic cells to specifically
recognize and destroy either cells infected with a
specific infective agent, the infective agent itself, a
15 tumor cell, or an autoimmune-generated cell. More
particularly, the invention relates to the production of
protein-tyrosine kinase chimeras capable of directing
cytotoxic T lymphocytes to specifically recognize and
lyse cells expressing HIV envelope proteins. The
20 invention therefore provides a therapy for diseases such
as AIDS (Acquired Immunodeficiency Syndrome) which are
caused by the HIV virus.

Background of the Invention

T cell recognition of antigen through the T cell
25 receptor is the basis of a range of immunological
phenomena. The T cells direct what is called cell-
mediated immunity. This involves the destruction by
cells of the immune system of foreign tissues or infected
cells. A variety of T cells exist, including "helper"
30 and "suppressor" cells, which modulate the immune
response, and cytotoxic (or "killer") cells, which can
kill abnormal cells directly.

A T cell that recognizes and binds a unique
antigen displayed on the surface of another cell becomes

- 2 -

activated; it can then multiply, and if it is a cytotoxic cell, it can kill the bound cell.

Autoimmune disease is characterized by production of either antibodies that react with host tissue or
5 immune effector T cells that are autoreactive. In some instances, autoantibodies may arise by a normal T- and B-cell response activated by foreign substances or organisms that contain antigens that cross react with similar compounds in body tissues. Examples of
10 clinically relevant autoantibodies are antibodies against acetylcholine receptors in myasthenia gravis; and anti-DNA, anti-erythrocyte, and anti-platelet antibodies in systemic lupus erythematosus.

HIV and Immunopathogenesis

15 In 1984 HIV was shown to be the etiologic agent of AIDS. Since that time the definition of AIDS has been revised a number of times with regard to what criteria should be included in the diagnosis. However, despite the fluctuation in diagnostic parameters, the simple
20 common denominator of AIDS is the infection with HIV and subsequent development of persistent constitutional symptoms and AIDS defining diseases such as a secondary infections, neoplasms, and neurologic disease.
Harrison's Principles of Internal Medicine, 12th ed.,
25 McGraw Hill (1991).

HIV is a human retrovirus of the lentivirus group. The four recognized human retroviruses belong to two distinct groups: the human T lymphotropic (or leukemia) retroviruses, HTLV-1 and HTLV-2, and the human
30 immunodeficiency viruses, HIV-1 and HIV-2. The former are transforming viruses whereas the latter are cytopathic viruses.

- 3 -

HIV-1 has been identified as the most common cause of AIDS throughout the world. Sequence homology between HIV-2 and HIV-1 is about 40% with HIV-2 being more closely related to some members of a group of simian immunodeficiency viruses (SIV). See Curran, J. et al., Science, 329:1357-1359 (1985); Weiss, R. et al., Nature, 324:572-575 (1986).

HIV has the usual retroviral genes (env, gag, and pol) as well as six extra genes involved in the replication and other biologic activities of the virus. As stated previously, the common denominator of AIDS is a profound immunosuppression, predominantly of cell-mediated immunity. This immune suppression leads to a variety of opportunistic diseases, particularly certain infections and neoplasms.

The main cause of the immune defect in AIDS, has been identified as a quantitative and qualitative deficiency in the subset of thymus-derived (T) lymphocytes, the T4 population. This subset of cells is defined phenotypically by the presence of the CD4 surface molecule, which has been demonstrated to be the cellular receptor for HIV. Dalglish et al., Nature, 312:763 (1984). Although the T4 cell is the major cell type infected with HIV, essentially any human cell that expresses the CD4 molecule on its surface is capable of binding to and being infected with HIV.

Traditionally, CD4⁺ T cells have been assigned the role of helper/inducer, indicating their function in providing an activating signal to B cells, or inducing T lymphocytes bearing the reciprocal CD8 marker to become cytotoxic/suppressor cells. Reinherz and Schlossman, Cell, 19:821-827 (1980); Goldstein et al., Immunol. Rev., 68:5-42, (1982).

HIV binds specifically and with high affinity, via a stretch of amino acids in the viral envelope (gp120),

- 4 -

to a portion of the V1 region of the CD4 molecule located near its N-terminus. Following binding, the virus fuses with the target cell membrane and is internalized. Once internalized it uses the enzyme reverse transcriptase to
5 transcribe its genomic RNA to DNA, which is integrated into the cellular DNA where it exists for the life of the cell as a "provirus."

The provirus may remain latent or be activated to transcribe mRNA and genomic RNA, leading to protein
10 synthesis, assembly, new virion formation, and budding of virus from the cell surface. Although the precise mechanism by which the virus induces cell death has not been established, it is felt that the major mechanism is massive viral budding from the cell surface, leading to
15 disruption of the plasma membrane and resulting osmotic disequilibrium.

During the course of the infection, the host organism develops antibodies against viral proteins, including the major envelope glycoproteins gp120 and
20 gp41. Despite this humoral immunity, the disease progresses, resulting in a lethal immunosuppression characterized by multiple opportunistic infections, parasitemia, dementia and death. The failure of the host anti-viral antibodies to arrest the progression of the
25 disease represents one of the most vexing and alarming aspects of the infection, and augurs poorly for vaccination efforts based upon conventional approaches.

Two factors may play a role in the efficacy of the humoral response to immunodeficiency viruses. First,
30 like other RNA viruses (and like retroviruses in particular), the immunodeficiency viruses show a high mutation rate in response to host immune surveillance. Second, the envelope glycoproteins themselves are heavily glycosylated molecules presenting few epitopes suitable
35 for high affinity antibody binding. The poorly antigenic

- 5 -

target which the viral envelope presents, allows the host little opportunity for restricting viral infection by specific antibody production.

Cells infected by the HIV virus express the gp120 glycoprotein on their surface. Gp120 mediates fusion events among CD4⁺ cells via a reaction similar to that by which the virus enters the uninfected cells, leading to the formation of short-lived multinucleated giant cells. Syncytium formation is dependent on a direct interaction of the gp120 envelope glycoprotein with the CD4 protein. Dalglish et al., supra; Klatzman, D. et al., Nature, 312:763 (1984); McDougal, J.S. et al., Science, 231:382 (1986); Sodroski, J. et al., Nature, 322:470 (1986); Lifson, J.D. et al., Nature, 323:725 (1986); Sodroski, J. et al., Nature, 321:412 (1986).

Evidence that the CD4-gp120 binding is responsible for viral infection of cells bearing the CD4 antigen includes the finding that a specific complex is formed between gp120 and CD4. McDougal et al., supra. Other investigators have shown that the cell lines, which were noninfective for HIV, were converted to infectable cell lines following transfection and expression of the human CD4 cDNA gene. Maddon et al., Cell, 46:333-348 (1986).

Therapeutic programs based on soluble CD4 as a passive agent to interfere with viral adsorption and syncytium-mediated cellular transmission have been proposed and successfully demonstrated in vitro by a number of groups (Deen et al., Nature, 332:82-84 (1988); Fisher et al., Nature, 331:76-78 (1988); Hussey et al., Nature 331:78-81 (1988); Smith et al., Science, 238:1704-1707 (1987); Trauneker et al., Nature, 331:84-86 (1988)); and CD4 immunoglobulin fusion proteins with extended halflives and modest biological activity have subsequently been developed (Capon et al., Nature, 337:525-531 (1989); Trauneker et al. Nature, 339, 68-70

- 6 -

(1989); Byrn et al., Nature, 344:667-670 (1990); Zettlmeissl et al., DNA Cell Biol. 9:347-353 (1990)). Although CD4 immunotoxin conjugates or fusion proteins show potent cytotoxicity for infected cells in vitro (Chaudhary et al., Nature, 335:369-372 (1988); Till et al., Science, 242:1166-1168 (1988)), the latency of the immunodeficiency syndrome makes it unlikely that any single-treatment therapy will be effective in eliminating viral burden, and the antigenicity of foreign fusion proteins is likely to limit their acceptability in treatments requiring repetitive dosing. Trials with monkeys affected with SIV have shown that soluble CD4, if administered to animals without marked CD4 cytopenia, can reduce SIV titer and improve in vitro measures of myeloid potential (Watanabe et al., Nature, 337:267-270 (1989)). However a prompt viral reemergence was observed after treatment was discontinued, suggesting that lifelong administration might be necessary to prevent progressive immune system debilitation.

20 Cell Surface Receptor-Associated Protein-Tyrosine Kinases

The initial impetus for engagement of cellular effector programs in the immune system is often cell recognition of clustered ligands. Among the receptors known to transmit activating signals upon aggregation are the B cell and T cell antigen receptors (DeFranco, 1992, Eur. J. Biochem. 210:381-388; Weiss, 1991, Annu. Rev. Genet. 25:487-510), members of the IgG and IgE Fc receptor families (Fanger et al., 1989, Immunol. Today 10:92-99; Ravetch and Kinet, 1991, Annu. Rev. Immunol. 9:457-492) and a number of accessory receptors, including CD2, CD4, CD8 and CD28 in T cells (Yokoyama and Shevach, 1989, Year Immunol. 4:110-146), CD19, CD20, CD21 and CD40 in B cells (Clark and Ledbetter, 1989, Adv. Cancer Res.

- 7 -

52:81-149), and CD44, CD45 and CD58 in monocytes (Webb et al., 1990, Science 249:1295-1297). In addition, a large number of phospholipid linked proteins promote cellular activation in an antigen receptor-dependent manner when
5 crosslinked on the surface of T cells (Balk and Terhorst, 1989, Immunol. Ser. 45:411-416; Kroczeck et al., 1986, Nature 322:181-184; Yeh et al., 1987, J. Immunol. 138:91-97; Yokoyama and Shevach, 1989, Year Immunol. 4:110-146).

At present it is not clear how a simple physical
10 event, aggregation, results in a clearly distinguished physiological signal. Engagement of cellular effector programs mediated by the T cell and B cell antigen receptors, and various forms of Fc receptor, can be mimicked by crosslinking of chimeric proteins bearing the
15 intracellular domains of individual chains of the receptor complexes (Irving and Weiss, 1991, Cell 64:891-901; Kolanus et al., 1992, EMBO J. 11:4861-4868; Letourneur and Klausner, 1991, Proc. Natl. Acad. Sci. USA 88:8905-8909; Letourneur and Klausner, 1992, Science
20 255:79-82; Romeo and Seed, 1991, Cell 64:1037-1046; Wegener et al., 1992, Cell 68:83-95). The minimal effective trigger element appears to require a phylogenetically conserved (Reth, 1989, Nature 338:383-384) peptide sequence containing two tyrosine residues
25 separated by 10 or 11 residues and embedded in a hydrophilic, typically acidic context (Romeo et al., 1992, Cell 68:889-897; Irving et al., 1993, J. Exp. Med. 177, 1093-1103). Clustering of receptors bearing this element initiates an activation cascade for which protein
30 tyrosine kinase (PTK) activity appears to be essential; PTK inhibitors block both early events in B and T cell activation such as calcium mobilization and the later sequelae of cytokine release and cellular proliferation (June et al., 1990, J. Immunol. 144:1591-1599; Lane et
35 al., 1991, J. Immunol. 146:715-722; Mustelin et al.,

- 8 -

1990, Science 247:1584-1587; Stanley et al., 1990, J. Immunol. 145:2189-2198). Although the more distal consequences of receptor activation differ according to cell type, the early events are strikingly similar among 5 cells from disparate hematopoietic lineages. For example the rapid increases in PTK activity observed following crosslinking of the B cell antigen receptor (Gold et al., 1990, Nature 345:810-813; Campbell and Sefton, 1990, EMBO J. 9:2125-2131), the T cell antigen receptor (June, C.H., 10 et al. 1990, Proc. Natl. Acad. Sci. USA 87:7722-7726; June, C.H., et al., 1990, J. Immunol. 144:1591-1599) and the high affinity IgE receptor (Eiseman and Bolen, 1992, Nature 355:78-80; Li et al., 1992, Mol. Cell. Biol. 12:3176-3182) all have among their early phosphorylation 15 targets the γ isoform of phosphatidylinositol-specific phospholipase C (Carter et al., 1991, Proc. Natl. Acad. Sci. USA 88:2745-2749; Li et al., 1992, Mol. Cell Biol. 12:3176-3182; Park et al., 1991, J. Biol. Chem. 266:24237-24240; Park et al., 1991, Proc. Natl. Acad. 20 Sci. USA 88:5453-5456; Secrist et al., 1991, J. Biol. Chem. 266:12135-12139; Weiss et al., 1991, Annu. Rev. Genet. 25:487-510), which is directly activated by tyrosine phosphorylation (Nishibe et al., 1990, Science 250:1253-1256).

25 The PTK activities known thus far to associate with cell surface receptors fall in two classes: those belonging to the family of Src proto-oncogene-related kinases and those related to the recently characterized Syk kinase. Among the former, the Fyn kinase has been 30 shown to associate with the T cell receptor (Gassmann et al., 1992, Eur. J. Immunol. 22:283-286; Samelson et al., 1990, Proc. Natl. Acad. Sci. USA 87:4358-4362), the Lyn, Fyn, Blk and Lck kinases have been reported to associate with the B cell IgM receptor, (Burkhardt et al., 1991, 35 Proc. Natl. Acad. Sci. USA 88:7410-7414; Campbell and

- 9 -

Sefton, 1992, Mol. Cell. Biol. 12:2315-2321; Yamanashi et al., 1991, Science 251:192-194), and the Lyn and Yes kinases have been shown to associate with the high affinity IgE receptor (Eiseman and Bolen, 1992, Nature 5 355:78-80; Hutchcroft et al., 1992, Proc. Natl. Acad. Sci. USA 89:9107-9111; Hutchcroft, J.E., et al., 1992, J. Biol. Chem. 267:8613-8619). The mechanism of the observed association has not been established in detail, but preliminary data suggest that the intracellular 10 domains of receptor complex chains may physically associate with Src family kinases (Clark et al., 1992, Science 258:123-126; Timson Gauen et al., 1992, Mol. Cell. Biol. 12:5438-5446). At present it is not clear whether these associations are direct or indirect.

15 To date, the most compelling evidence for the importance of Src family kinases in cell activation has been developed from the study of the Fyn and Lck kinases in T cells. Overexpression of Fyn in transgenic mice leads to an antigen hyperresponsive phenotype in the 20 resulting T cells, while overexpression of a catalytically inactive form blocks T cell receptor mediated proliferation (Cooke et al., 1991, Cell 65:281-291). Thymic T cells isolated from mutant mice lacking Fyn kinase activity show a profound defect in the ability 25 to mount a proliferative response in response to treatment with a combination of phorbol ester plus either anti-CD3 antibody or Concanavalin A (Appleby et al., 1992, Cell 70:751-763; Stein et al., 1992, Cell 70:741-750). Splenic T cells isolated from such mice show a 30 less severe, but substantial, attenuation of the cell activation response (Appleby et al., 1992, Cell 70:751-763; Stein et al., 1992, Cell 70:741-750).

In T cells the Lck kinase associates indirectly with the TCR through the CD4 and CD8 coreceptors (Rudd et 35 al., 1988, Proc. Natl. Acad. Sci. USA 85:5190-5194; Shaw

- 10 -

et al., 1989, Cell 59:627-636; Turner et al., 1990, Cell 60:755-765; Veillette et al., 1988, Cell 55:301-308). Overexpression of Lck in an antigen-responsive cell line potentiates receptor sensitivity in similar fashion to
5 that seen with Fyn (Abraham and Veillette, 1990, Mol. Cell. Biol. 10:5197-5206; Davidson et al., 1992, J. Exp. Med. 175:1483-1492; Luo and Sefton, 1992, Mol. Cell. Biol. 12:4724-4732). In a CD4-dependent murine T cell hybridoma model, reconstitution of antigen-specific
10 helper function could be achieved only with CD4 molecules which were capable of interacting with Lck (Glaichenhaus et al., 1991, Cell 64:511-520).

However the strongest evidence for the direct participation of the Lck kinase in antigen
15 receptor-mediated signalling comes from studies of mutant cell lines which lack Lck. Two such lines have been studied, one derived from the Jurkat human T cell leukemia line (Goldsmith and Weiss, 1987, Proc. Natl. Acad. Sci. USA 84:6879-6883; Straus and Weiss, 1992, Cell
20 70:585-593) and the other from the murine cytotoxic T cell clone CTLL-2 (Karnitz et al., 1992, Mol. Cell. Biol. 12:4521-4530). Both Lck-negative mutant lines are defective in TCR mediated signalling, and complementation of either mutant line by transfection with an Lck
25 expression plasmid restores responsiveness to TCR crosslinking stimuli (Karnitz et al., 1992, Mol. Cell. Biol. 12:4521-4530; Straus and Weiss, 1992, Cell 70:585-593).

Recently members of a new family of tyrosine
30 kinases, initially represented by the closely related or identical kinases Syk (Taniguchi et al., 1991, J. Biol. Chem. 266:15790-15796) and PTK 72 (Zioncheck et al., 1986, J. Biol. Chem. 261:15637-15643; Zioncheck et al., 1988, J. Biol. Chem. 263:19195-19202), have been shown
35 to associate with cell surface receptors. Although PTK

- 11 -

72 and Syk have not been definitively proven to be identical, they share a common tissue distribution (thymus and spleen), molecular mass, and lability to proteolysis. PTK 72 has been shown to associate with the B cell IgM receptor (Hutchcroft et al., 1992, Proc. Natl. Acad. Sci. USA 89:9107-9111; Hutchcroft, J.E., et al., 1992, J. Biol. Chem. 267:8613-8619) and to be phosphorylated upon crosslinking of the receptor with anti-IgM (Hutchcroft et al., 1991, J. Biol. Chem. 266:14846-14849). A concomitant activation of the enzyme, as measured by both autophosphorylation and phosphorylation of an exogenous protein fragment, was demonstrated following surface IgM crosslinking (Hutchcroft et al., 1992, Proc. Natl. Acad. Sci. USA 89:9107-9111; Hutchcroft, J.E., et al., 1992, J. Biol. Chem. 267:8613-8619). PTK 72 is also found associated with the high affinity IgE receptor in a rat basophilic leukemia cell line (Hutchcroft et al., 1992, Proc. Natl. Acad. Sci. USA 89:9107-9111; Hutchcroft, J.E., et al., 1992, J. Biol. Chem. 267:8613-8619).

A second member of the Syk family, ZAP-70, has been shown to be a PTK associating with the zeta chain of the T cell receptor following receptor crosslinking (Chan et al., 1991, Proc. Natl. Acad. Sci. USA 88:9166-9170). Although expression in COS cells of ZAP-70, Fyn or Lck leads to modest increases in total cell tyrosine phosphate, coexpression of ZAP-70 and either Lck or Fyn leads to a dramatic increase in net tyrosine phosphorylation (Chan et al., 1992, Cell 71:649-662). If a CD8-zeta chain chimera is also present, the chimera becomes phosphorylated and ZAP-70 is found associated with it (Chan et al., 1992, Cell 71:649-662). At present it is not clear whether ZAP-70 activates the Src family kinases and/or vice versa, nor why coexpression of kinases in COS cells should lead to an apparent

- 12 -

constitutive activation. Nonetheless the active association of ZAP-70 with crosslinked TCR suggests a role for this PTK in the propagation of the receptor response.

5 Unlike the Src family kinases, Syk and ZAP-70 bear two SH2 domains and no N-terminal myristoylation site (Taniguchi et al., 1991, J. Biol. Chem. 266:15790-15796; Chan et al., 1992, Cell 71:649-662). A natural expectation for the mechanism of kinase-receptor
10 association is that the two SH2 domains bind the two tyrosines of the antigen receptor trigger motifs once they are phosphorylated. However, at present this is remains merely a hypothesis.

Summary of the Invention

15 The present invention demonstrates the feasibility of creating chimeras between the intracellular domain of a protein-tyrosine kinase molecule and an extracellular domain which is capable of fulfilling the task of target recognition. In particular, clustering of chimeras
20 bearing Syk or ZAP-70 kinase sequences triggers calcium mobilization. Aggregation of Syk chimera alone, or coaggregation of chimeras bearing Fyn or Lck and ZAP-70 kinases, suffices to initiate cytolytic effector function. Such effector function facilitates the
25 specific recognition and destruction of undesirable target cells, for example, pathogens, pathogen-infected cells, tumor cells, or autoimmune cells.

Any number of useful chimeric molecules according to the invention may be constructed. For example, the
30 formation of chimeras consisting of the intracellular portion of a protein-tyrosine kinase joined to the extracellular portion of a suitably engineered antibody molecule allows the target recognition potential of an immune system cell to be specifically redirected to the

- 13 -

antigen recognized by the extracellular antibody portion. Thus with an antibody portion capable of recognizing some determinant on the surface of a pathogen, immune system cells armed with the chimera would respond to the
5 presence of the pathogen with the effector program appropriate to their lineage, e.g., helper T lymphocytes would respond by cytotoxic activity against the target, and B lymphocytes would be activated to synthesize antibody. Macrophages and granulocytes would carry out
10 their effector programs, including cytokine release, phagocytosis, and reactive oxygen generation. Similarly, with an antibody portion capable of recognizing tumor cells, the immune system response to the tumor would be beneficially elevated. With an antibody capable of
15 recognizing immune cells having an inappropriate reactivity with self determinants, the autoreactive cells could be selectively targeted for destruction.

Although these examples draw on the use of antibody chimeras as a convenient expository tool, the
20 invention is not limited in scope to antibody chimeras, and indeed, the use of specific nonantibody extracellular domains may have important advantages. For example with an extracellular portion that is the receptor for a virus, bacterium, or parasite, cells armed with the
25 chimeras would specifically target cells expressing the viral, bacterial or parasitic determinants. The advantage of this approach over the use of antibodies is that the native receptor for pathogen may have uniquely high selectivity or affinity for the pathogen, allowing a
30 greater degree of precision in the resulting immune response. Similarly, to delete immune system cells which inappropriately react with a self antigen, it may suffice to join the antigen (either as an intact protein, in the case of B cell depletion therapies, or as MHC complex, in
35 the case of T cell depletion therapies) to intracellular

- 14 -

protein-tyrosine kinase chains, and thereby affect the specific targeting of the cells inappropriately responding to self determinants.

Another use of the chimeras is the control of cell
5 populations in vivo subsequent to other forms of genetic engineering. For example, the use of tumor infiltrating lymphocytes or natural killer cells to carry cytotoxic principles to the site of tumors has been proposed. The present invention provides a convenient means to regulate
10 the numbers and activity of such lymphocytes and cells without removing them from the body of the patient for amplification in vitro. Thus, because the intracellular domains of the chimeric receptors mediate the proliferative responses of the cells, the coordination of
15 the extracellular domains by a variety of aggregating stimuli specific for the extracellular domains (e.g., an antibody specific for the extracellular domain) will result in proliferation of the cells bearing the chimeras.

20 Although the specific embodiments of the present invention comprise chimeras between the Syk or Syk and Src families of protein-tyrosine kinases, any tyrosine kinase having a similar function to these molecules could be used for the purposes disclosed here. The
25 distinguishing features of desirable immune cell trigger molecules comprise the ability to be expressed autonomously, the ability to be fused to an extracellular domain (directly or indirectly through a transmembrane domain) such that the resultant chimera is present on the
30 surface of a therapeutic cell, and the ability to initiate cellular effector programs upon aggregation secondary to encounter with a target ligand.

At present the most convenient method for delivery of the chimeras to immune system cells is through some
35 form of genetic therapy. However reconstituting immune

- 15 -

system cells with chimeric receptors by mixture of the cells with suitably solubilized purified chimeric protein would also result in the formation of an engineered cell population capable of responding to the targets

5 recognized by the extracellular domain of the chimeras. Similar approaches have been used, for example, to introduce the intact HIV receptor, CD4, into erythrocytes for therapeutic purposes. In this case the engineered cell population would not be capable of self renewal.

10 The present invention relates to functional simplified protein-tyrosine kinase chimeras which are capable of redirecting immune system function. More particularly, it relates to the regulation of lymphocytes, macrophages, natural killer cells or
15 granulocytes by the expression in said cells of chimeras which cause the cells to respond to targets recognized by the chimeras. The invention also relates to a method of directing cellular response to an infective agent, a tumor or cancerous cell, or an autoimmune generated cell.

20 The method for directing the cellular response in a mammal comprises administering an effective amount of therapeutic cells to said mammal, said cells being capable of recognizing and destroying said infective agent, tumor, cancer cell, or autoimmune generated cell.

25 In another embodiment, the method of directing cellular response to an infective agent comprises administering therapeutic cells capable of recognizing and destroying said agent, wherein the agent is a specific virus, bacteria, protozoa, or fungi. Even more
30 specifically, the method is directed against agents such as HIV and Pneumocystis carinii.

To treat an HIV infection, an effective amount of chimeric-receptor expression cytotoxic T lymphocytes are administered to a patient; the lymphocytes are capable of

- 16 -

specifically recognizing and lysing cells infected with HIV as well as circulating virus.

Thus, in one embodiment, there is provided according to the invention a method for directing
5 cellular response to HIV infected cells, comprising administering to a patient an effective amount of cytotoxic T lymphocytes which are capable of specifically recognizing and lysing cells infected with HIV.

In yet another embodiment is provided the chimeric
10 receptor proteins which direct the cytotoxic T lymphocytes to recognize and lyse the HIV infected cell. Yet another embodiment of the invention comprises host cells transformed with a vector comprising the chimeric receptors.

15 These and other non-limiting embodiments of the present invention will be apparent to those of skill from the following detailed description of the invention.

In the following detailed description, reference will be made to various methodologies known to those of
20 skill in the art of molecular biology and immunology. Publications and other materials setting forth such known methodologies to which reference is made are incorporated herein by reference in their entireties as though set forth in full.

25 Standard reference works setting forth the general principles of recombinant DNA technology include Watson, J.D. et al., Molecular Biology of the Gene, Volumes I and II, the Benjamin/Cummings Publishing Company, Inc., publisher, Menlo Park, CA (1987); Darnell, J.E. et al.,
30 Molecular Cell Biology, Scientific American Books, Inc., Publisher, New York, N.Y. (1986); Lewin, B.M., Genes II, John Wiley & Sons, publishers, New York, N.Y. (1985); Old, R.W., et al., Principles of Gene Manipulation: An Introduction to Genetic Engineering, 2d edition,
35 University of California Press, publisher, Berkeley, CA

- 17 -

(1981); Maniatis, T., et al., Molecular Cloning: A Laboratory Manual, 2nd Ed. Cold Spring Harbor Laboratory, publisher, Cold Spring Harbor, NY (1989); and Current Protocols in Molecular Biology, Ausubel et al., Wiley Press, New York, NY (1989).

DEFINITIONS

By "cloning" is meant the use of in vitro recombination techniques to insert a particular gene or other DNA sequence into a vector molecule. In order to
10 successfully clone a desired gene, it is necessary to employ methods for generating DNA fragments for joining the fragments to vector molecules, for introducing the composite DNA molecule into a host cell in which it can replicate, and for selecting the clone having the target
15 gene from amongst the recipient host cells.

By "cDNA" is meant complementary or copy DNA produced from an RNA template by the action of RNA-dependent DNA polymerase (reverse transcriptase). Thus a "cDNA clone" means a duplex DNA sequence complementary to
20 an RNA molecule of interest, carried in a cloning vector.

By "cDNA library" is meant a collection of recombinant DNA molecules containing cDNA inserts which comprise DNA copies of mRNA being expressed by the cell at the time the cDNA library was made. Such a cDNA
25 library may be prepared by methods known to those of skill, and described, for example, in Maniatis et al., Molecular Cloning: A Laboratory Manual, supra.

Generally, RNA is first isolated from the cells of an organism from whose genome it is desired to clone a
30 particular gene. Preferred for the purpose of the present invention are mammalian, and particularly human, lymphocytic cell lines. A presently preferred vector for this purpose is the vaccinia virus WR strain.

- 18 -

By "vector" is meant a DNA molecule, derived, e.g., from a plasmid, bacteriophage, or mammalian or insect virus, into which fragments of DNA may be inserted or cloned. A vector will contain one or more unique
5 restriction sites and may be capable of autonomous replication in a defined host or vehicle organism such that the cloned sequence is reproducible. Thus, by "DNA expression vector" is meant any autonomous element capable of directing the synthesis of a recombinant
10 peptide. Such DNA expression vectors include bacterial plasmids and phages and mammalian and insect plasmids and viruses.

By "substantially pure" is meant a compound, e.g., a protein, a polypeptide, or an antibody, that is
15 substantially free of the components that naturally accompany it. Generally, a compound is substantially pure when at least 60%, more preferably at least 75%, and most preferably at least 90% of the total material in a sample is the compound of interest. Purity can be
20 measured by any appropriate method, e.g., column chromatography, polyacrylamide gel electrophoresis, or HPLC analysis. In the context of a nucleic acid, "substantially pure" means a nucleic acid sequence, segment, or fragment that is not immediately contiguous
25 with (i.e., covalently linked to) both of the coding sequences with which it is immediately contiguous (i.e., one at the 5' end and one at the 3' end) in the naturally occurring genome of the organism from which the DNA of the invention is derived.

30 By "functional derivative" is meant the "fragments," "variants," "analogues," or "chemical derivatives" of a molecule. A "fragment" of a molecule, such as any of the cDNA sequences of the present invention, is meant to refer to any nucleotide subset of
35 the molecule. A "variant" of such molecule is meant to

- 19 -

refer to a naturally occurring molecule substantially similar to either the entire molecule, or a fragment thereof. An "analog" of a molecule is meant to refer to a non-natural molecule substantially similar to either
5 the entire molecule or a fragment thereof. A molecule is said to be "substantially similar" to another molecule if the sequence of amino acids in both molecules is substantially the same. Substantially similar amino acid molecules will possess a similar biological activity.
10 Thus, provided that two molecules possess a similar activity, they are considered variants as that term is used herein even if one of the molecules contains additional or fewer amino acid residues not found in the other, or if the sequence of amino acid residues is not
15 identical. As used herein, a molecule is said to be a "chemical derivative" of another molecule when it contains additional chemical moieties not normally a part of the molecule. Such moieties may improve the molecule's solubility, absorption, biological half life,
20 etc. The moieties may alternatively decrease the toxicity of the molecule, eliminate or attenuate any undesirable side effect of the molecule, etc. Moieties capable of mediating such effects are disclosed, for example, in Remington's Pharmaceutical Sciences, 16th
25 et., Mack Publishing Co., Easton, Penn. (1980).

Similarly, a "functional derivative" of a receptor chimera gene of the present invention is meant to include "fragments," "variants," or "analogues" of the gene, which may be "substantially similar" in nucleotide
30 sequence, and which encode a molecule possessing similar activity to a protein-tyrosine kinase chimera.

Thus, as used herein, a protein-tyrosine kinase chimera protein is also meant to include any functional derivative, fragments, variants, analogues, or chemical
35 derivatives which may be substantially similar to the

- 20 -

"wild-type" chimera and which possess similar activity (i.e., most preferably, 90%, more preferably, 70%, preferably 40%, or at least 10% of the wild-type receptor chimera's activity). The activity of a functional
5 chimeric receptor derivative includes specific binding (with its extracellular portion) to a targeted agent or cell and resultant destruction (directed by its intracellular portion) of that agent or cell; such activity may be tested, e.g., using any of the assays
10 described herein.

A DNA sequence encoding the chimera of the present invention, or its functional derivatives, may be recombined with vector DNA in accordance with conventional techniques, including blunt-ended or
15 staggered-ended termini for ligation, restriction enzyme digestion to provide appropriate termini, filling in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and ligation with appropriate ligases. Techniques for such manipulations
20 are disclosed by Maniatis, T., et al., supra, and are well known in the art.

A nucleic acid molecule, such as DNA, is said to be "capable of expressing" a polypeptide if it contains nucleotide sequences which contain transcriptional and
25 translational regulatory information and such sequences are "operably linked" to nucleotide sequences which encode the polypeptide. An operable linkage is a linkage in which the regulatory DNA sequences and the DNA sequence sought to be expressed are connected in such a
30 way as to permit gene expression. The precise nature of the regulatory regions needed for gene expression may vary from organism to organism, but shall in general include a promoter region which, in prokaryotes, contains both the promoter (which directs the initiation of RNA
35 transcription) as well as the DNA sequences which, when

- 21 -

transcribed into RNA, will signal the initiation of protein synthesis. Such regions will normally include those 5'-non-coding sequences involved with initiation of transcription and translation, such as the TATA box, 5 capping sequence, CAAT sequence, and the like.

If desired, the non-coding region 3' to the gene sequence coding for the protein may be obtained by the above-described methods. This region may be retained for its transcriptional termination regulatory sequences, 10 such as termination and polyadenylation. Thus, by retaining the 3'-region naturally contiguous to the DNA sequence coding for the protein, the transcriptional termination signals may be provided. Where the transcriptional termination signals are not 15 satisfactorily functional in the expression host cell, then a 3' region functional in the host cell may be substituted.

Two DNA sequences (such as a promoter region sequence and a protein-tyrosine kinase chimera-encoding 20 sequence) are said to be operably linked if the nature of the linkage between the two DNA sequences does not (1) result in the introduction of a frame-shift mutation, (2) interfere with the ability of the promoter region sequence to direct the transcription of the receptor 25 chimera gene sequence, or (3) interfere with the ability of the receptor chimera gene sequence to be transcribed by the promoter region sequence. A promoter region would be operably linked to a DNA sequence if the promoter were capable of effecting transcription of that DNA sequence. 30 Thus, to express the protein, transcriptional and translational signals recognized by an appropriate host are necessary.

The present invention encompasses the expression of a protein-tyrosine kinase chimera protein (or a 35 functional derivative thereof) in either prokaryotic or

- 22 -

eukaryotic cells, although eukaryotic (and, particularly, human lymphocyte) expression is preferred.

Antibodies according to the present invention may be prepared by any of a variety of methods. For example, 5 cells expressing the receptor chimera protein, or a functional derivative thereof, can be administered to an animal in order to induce the production of sera containing polyclonal antibodies that are capable of binding the chimera.

10 In a preferred method, antibodies according to the present invention are monoclonal antibodies. Such monoclonal antibodies can be prepared using hybridoma technology (Kohler et al., Nature 256:495 (1975); Kohler et al., Eur. J. Immunol. 6:511 (1976); Kohler et al., 15 Eur. J. Immunol. 6:292 (1976); Hammerling et al., In: Monoclonal Antibodies and T-Cell Hybridomas, Elsevier, N.Y., pp. 563-684 (1981)). In general, such procedures involve immunizing an animal with the chimera antigen. The splenocytes of such animals are extracted and fused 20 with a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the present invention. After fusion, the resulting hybridoma cells are selectively maintained in HAT medium, and then cloned by limiting dilution as described by Wands, J.R., et al. 25 (Gastroenterology 80:225-232 (1981)). The hybridoma cells obtained through such a selection are then assayed to identify clones which secrete antibodies capable of binding the chimera.

Antibodies according to the present invention also 30 may be polyclonal, or, preferably, region specific polyclonal antibodies.

Antibodies against the chimera according to the present invention may be used to monitor the amount of chimeric receptor (or chimeric receptor-bearing cells) in 35 a patient. Such antibodies are well suited for use in

- 23 -

standard immunodiagnostic assay known in the art, including such immunometric or "sandwich" assays as the forward sandwich, reverse sandwich, and simultaneous sandwich assays. The antibodies may be used in any
5 number of combinations as may be determined by those of skill without undue experimentation to effect immunoassays of acceptable specificity, sensitivity, and accuracy.

Standard reference works setting forth general
10 principles of immunology include Roitt, I., Essential Immunology, Sixth Ed., Blackwell Scientific Publications, Publisher, Oxford (1988); Kimball, J. W., Introduction to Immunology, Second Ed., Macmillan Publishing Co., Publisher, New York (1986); Roitt, I., et al.,
15 Immunology, Gower Medical Publishing Ltd., Publisher, London, (1985); Campbell, A. , "Monoclonal Antibody Technology," in, Burdon, R., et al., eds., Laboratory Techniques in Biochemistry and Molecular Biology, Volume 13, Elsevier, Publisher, Amsterdam (1984); Klein, J.,
20 Immunology: The Science of Self-Nonself Discrimination, John Wiley & Sons, Publisher, New York (1982); and Kennett, R., et al., eds., Monoclonal Antibodies, Hybridoma: A New Dimension In Biological Analyses, Plenum Press, Publisher, New York (1980).

25 By "detecting" it is intended to include determining the presence or absence of a substance or quantifying the amount of a substance. The term thus refers to the use of the materials, compositions, and methods of the present invention for qualitative and
30 quantitative determinations.

The isolation of other hybridomas secreting monoclonal antibodies of the same specificity as those described herein can be accomplished by the technique of anti-idiotypic screening (Potocmjak, et al., Science
35 215:1637 (1982)). Briefly, an anti-idiotypic antibody is

- 24 -

an antibody which recognizes unique determinants present on the antibody produced by the clone of interest. The anti-idiotypic antibody is prepared by immunizing an animal of the same strain used as the source of the
5 monoclonal antibody with the monoclonal antibody of interest. The immunized animal will recognize and respond to the idiotypic determinants of the immunizing antibody by producing antibody to these idiotypic determinants (anti-idiotypic antibody).

10 For replication, the hybrid cells may be cultivated both in vitro and in vivo. High in vivo production makes this the presently preferred method of culture. Briefly, cells from the individual hybrid strains are injected intraperitoneally into pristane-
15 primed BALB/c mice to produce ascites fluid containing high concentrations of the desired monoclonal antibodies. Monoclonal antibodies of isotype IgM or IgG may be purified from cultured supernatants using column chromatography methods well known to those of skill in
20 the art.

Antibodies according to the present invention are particularly suited for use in immunoassays wherein they may be utilized in liquid phase or bound to a solid phase carrier. In addition, the antibodies in these
25 immunoassays can be detectably labeled in various ways.

There are many different labels and methods of labeling known in the art. Examples of the types of labels which can be used in the present invention include, but are not limited to, enzymes, radioisotopes,
30 fluorescent compounds, chemiluminescent compounds, bioluminescent compounds and metal chelates. Those of ordinary skill in the art will know of other suitable labels for binding to antibodies, or will be able to ascertain the same by the use of routine experimentation.
35 Furthermore, the binding of these labels to antibodies

- 25 -

can be accomplished using standard techniques commonly known to those of ordinary skill in the art.

One of the ways in which antibodies according to the present invention can be detectably labeled is by
5 linking the antibody to an enzyme. This enzyme, in turn, when later exposed to its substrate, will react with the substrate in such a manner as to produce a chemical moiety which can be detected as, for example, by spectrophotometric or fluorometric means. Examples of
10 enzymes which can be used to detectably label antibodies include malate dehydrogenase, staphylococcal nuclease, delta-V-steroid isomerase, yeast alcohol dehydrogenase, alpha-glycerophosphate dehydrogenase, triose phosphate isomerase, biotinavidin peroxidase, horseradish
15 peroxidase, alkaline phosphatase, asparaginase, glucose oxidase, β -galactosidase, ribonuclease, urease, catalase, glucose-VI-phosphate dehydrogenase, glucoamylase and acetylcholine esterase.

The presence of detectably labeled antibodies also
20 can be detected by labeling the antibodies with a radioactive isotope which then can be determined by such means as the use of a gamma counter or a scintillation counter. Isotopes which are particularly useful for the purpose of the present invention are ^3H , ^{125}I , ^{32}P , ^{35}S ,
25 ^{14}C , ^{51}Cr , ^{36}Cl , ^{57}Co , ^{58}Co , ^{59}Fe and ^{75}Se .

It is also possible to detect the binding of detectably labeled antibodies by labeling the antibodies with a fluorescent compound. When a fluorescently labeled antibody is exposed to light of the proper
30 wavelength, its presence then can be detected due to the fluorescence of the dye. Among the most commonly used fluorescent labeling compounds are fluorescein, isothiocyanate, rhodamine, phycoerythrin, phycocyanin, allophycocyanin, o-phthaldehyde and fluorescamine.

- 26 -

The antibodies of the invention also can be detectably labeled using fluorescent emitting metals such as ^{152}Eu , or others of the lanthanide series. These metals can be attached to the antibody molecule using
5 such metal chelating groups as diethyl-
enteriaminepentaacetic acid (DTPA) or
ethylenediaminetetraacetic acid (EDTA).

Antibodies also can be detectably labeled by coupling them to a chemiluminescent compound. The
10 presence of the chemiluminescent-tagged antibody is then determined by detecting the presence of luminescence that arises during the course of the chemical reaction. Examples of particularly useful chemiluminescent labeling compounds are luminal, isoluminol, theromatic acridinium
15 ester, imidazole, acridinium salts, oxalate ester, and dioxetane.

Likewise, a bioluminescent compound may be used to label the antibodies according to the present invention. Bioluminescence is a type of chemiluminescence found in
20 biological systems in which a catalytic protein increases the efficiency of the chemiluminescent reaction. The presence of a bioluminescent antibody is determined by detecting the presence of luminescence. Important bioluminescent compounds for purposes of labeling include
25 luciferin, luciferase aequorin.

The antibodies and substantially purified antigen of the present invention are ideally suited for the preparation of a kit. Such a kit may comprise a carrier means being compartmentalized to receive in close
30 confinement therewith one or more container means such as vials, tubes and the like, each of said container means comprising the separate elements of the assay to be used.

The types of assays which can be incorporated in kit form are many, and include, for example, competitive
35 and non-competitive assays. Typical examples of assays

- 27 -

which can utilize the antibodies of the invention are radioimmunoassays (RIA), enzyme immunoassays (EIA), enzyme-linked immunosorbent assays (ELISA), and immunometric, or sandwich, immunoassays.

5 By the term "immunometric assay" or "sandwich immunoassay," it is meant to include simultaneous sandwich, forward sandwich and reverse sandwich immunoassays. These terms are well understood by those skilled in the art. Those of skill will also appreciate
10 that antibodies according to the present invention will be useful in other variations and forms of assays which are presently known or which may be developed in the future. These are intended to be included within the scope of the present invention.

15 In the preferred mode for performing the assays it is important that certain "blockers" be present in the incubation medium (usually added with the labeled soluble antibody). The "blockers" are added to assure that non-specific proteins, protease, or human antibodies to mouse
20 immunoglobulins present in the experimental sample do not cross-link or destroy the antibodies on the solid phase support, or the radiolabeled indicator antibody, to yield false positive or false negative results. The selection of "blockers" therefore adds substantially to the
25 specificity of the assays described in the present invention.

It has been found that a number of nonrelevant (i.e., nonspecific) antibodies of the same class or subclass (isotype) as those used in the assays (e.g.,
30 IgG₁, IgG_{2a}, IgM, etc.) can be used as "blockers." The concentration of the "blockers" (normally 1-100 µg/µl) is important, in order to maintain the proper sensitivity yet inhibit any unwanted interference by mutually occurring cross reactive proteins in human serum. In
35 addition, the buffer system containing the "blockers"

- 28 -

needs to be optimized. Preferred buffers are those based on weak organic acids, such as imidazole, HEPPS, MOPS, TES, ADA, ACES, HEPES, PIPES, TRIS, and the like, at physiological pH ranges. Somewhat less preferred buffers
5 are inorganic buffers such as phosphate, borate or carbonate. Finally, known protease inhibitors should be added (normally at 0.01-10 $\mu\text{g/ml}$) to the buffer which contains the "blockers."

There are many solid phase immunoabsorbents which
10 have been employed and which can be used in the present invention. Well known immunoabsorbents include glass, polystyrene, polypropylene, dextran, nylon and other materials, in the form of tubes, beads, and microtiter plates formed from or coated with such materials, and the
15 like. The immobilized antibodies can be either covalently or physically bound to the solid phase immunoabsorbent, by techniques such as covalent bonding via an amide or ester linkage, or by absorption. Those skilled in the art will know many other suitable solid
20 phase immunoabsorbents and methods for immobilizing antibodies thereon, or will be able to ascertain such, using no more than routine experimentation.

For in vivo, in vitro, or in situ diagnosis, labels such as radionuclides may be bound to antibodies
25 according to the present invention either directly or by using an intermediary functional group. An intermediary group which is often used to bind radioisotopes which exist as metallic cations to antibodies is diethylenetriaminepentaacetic acid (DTPA). Typical
30 examples of metallic cations which are bound in this manner are: $^{99\text{m}}\text{Tc}$, ^{123}I , ^{111}In , ^{131}I , ^{97}Ru , ^{67}Cu , ^{67}Ga and ^{68}Ga . The antibodies of the invention can also be labeled with non-radioactive isotopes for purposes of diagnosis. Elements which are particularly useful in this manner are
35 ^{157}Gd , ^{55}Mn , ^{162}Dy , ^{52}Cr and ^{56}Fe .

- 29 -

The antigen of the invention may be isolated in substantially pure form employing antibodies according to the present invention. Thus, an embodiment of the present invention provides for substantially pure
5 protein-tyrosine kinase chimera, said antigen characterized in that it is recognized by and binds to antibodies according to the present invention. In another embodiment, the present invention provides a method of isolating or purifying the chimeric receptor
10 antigen, by forming a complex of said antigen with one or more antibodies directed against the receptor chimera.

The substantially pure chimera antigens of the present invention may in turn be used to detect or measure antibody to the chimera in a sample, such as
15 serum or urine. Thus, one embodiment of the present invention comprises a method of detecting the presence or amount of antibody to protein-tyrosine kinase antigen in a sample, comprising contacting a sample containing an antibody to the chimeric antigen with detectably labeled
20 receptor chimera, and detecting said label. It will be appreciated that immunoreactive fractions and immunoreactive analogues of the chimera also may be used. By the term "immunoreactive fraction" is intended any portion of the chimeric antigen which demonstrates an
25 equivalent immune response to an antibody directed against the receptor chimera. By the term "immunoreactive analogue" is intended a protein which differs from the receptor chimera protein by one or more amino acids, but which demonstrates an equivalent
30 immunoreponse to an antibody of the invention.

By "specifically recognizes and binds" is meant that the antibody recognizes and binds the chimeric receptor polypeptide but does not substantially recognize and bind other unrelated molecules in a sample, e.g., a
35 biological sample.

- 30 -

By "autoimmune-generated cell" is meant cells producing antibodies that react with host tissue or immune effector T cells that are autoreactive; such cells include antibodies against acetylcholine receptors (leading, e.g., to myasthenia gravis) or anti-DNA, anti-erythrocyte, and anti-platelet autoantibodies (leading, e.g., to lupus erythematosus).

By "therapeutic cell" is meant a cell which has been transformed by a chimera of the invention so that it is capable of recognizing and destroying a specific infective agent, a cell infected by a specific agent, a tumor or cancerous cell, or an autoimmune-generated cell; preferably such therapeutic cells are cells of the hematopoietic system.

By a "target infective agent" is meant any infective agent (e.g., a virus, bacterium, protozoan, or fungus) which can be recognized by a chimeric receptor-bearing therapeutic cell. By a "target cell" is meant any host cell which can be recognized by a chimeric receptor-bearing therapeutic cell; target cells include, without limitation, host cells which are infected with a virus, bacterium, protozoan, or fungus as well as tumor or cancerous cells and autoimmune-generated cells.

By "extracellular" is meant having at least a portion of the molecule exposed at the cell surface. By "intracellular" is meant having at least a portion of the molecule exposed to the therapeutic cell's cytoplasm. By "transmembrane" is meant having at least a portion of the molecule spanning the plasma membrane. An "extracellular portion", an "intracellular portion" and a "transmembrane portion", as used herein, may include flanking amino acid sequences which extend into adjoining cellular compartments.

By "oligomerize" is meant to complex with other proteins to form dimers, trimers, tetramers, or other

- 31 -

higher order oligomers. Such oligomers may be homo-oligomers or hetero-oligomers. An "oligomerizing portion" is that region of a molecule which directs complex (i.e., oligomer) formation.

5 By "cytolytic" is meant to be capable of destroying a cell (e.g., a cell infected with a pathogen, a tumor or cancerous cell, or an autoimmune-generated) cell or to be capable of destroying an infective agent (e.g., a virus).

10 By "immunodeficiency virus" is meant a retrovirus that, in wild-type form, is capable of infecting T4 cells of a primate host and possesses a viral morphogenesis and morphology characteristic of the lentivirus subfamily. The term includes, without limitation, all variants of
15 HIV and SIV, including HIV-1, HIV-2, SIVmac, SIVagm, SIVmnd, SIVsmm, SIVman, SIVmand, and SIVcpz.

By "MHC-independent" is meant that the cellular cytolytic response does not require the presence of an MHC class II antigen on the surface of the targeted cell.

20 By a "functional cytolytic signal-transducing derivative" is meant a functional derivative (as defined above) which is capable of directing at least 10%, preferably 40%, more preferably 70%, or most preferably at least 90% of the biological activity of the wild type
25 molecule. As used herein, a "functional cytolytic signal-transducing derivative" may act by directly signaling the therapeutic cell to destroy a receptor-bound agent or cell (e.g., in the case of an intracellular chimeric receptor portion) or may act
30 indirectly by promoting oligomerization with cytolytic signal transducing proteins of the therapeutic cell (e.g., in the case of a transmembrane domain). Such derivatives may be tested for efficacy, e.g., using the in vitro assays described herein.

- 32 -

By a "functional HIV envelope-binding derivative" is meant a functional derivative (as defined above) which is capable of binding any HIV envelope protein. Functional derivatives may be identified using, e.g., the
5 in vitro assays described herein.

THERAPEUTIC ADMINISTRATION

The transformed cells of the present invention may be used for the therapy of a number of diseases. Current methods of administering such transformed cells involve
10 adoptive immunotherapy or cell-transfer therapy. These methods allow the return of the transformed immune-system cells to the bloodstream. Rosenberg, S.A., Scientific American, 62 (May 1990); Rosenberg et al., The New England Journal of Medicine, 323(9):570 (1990).

15 The pharmaceutical compositions of the invention may be administered to any animal which may experience the beneficial effects of the compounds of the invention. Foremost among such animals are humans, although the invention is not intended to be so limited.

20

Detailed Description

The drawings will first be described.

Description of the Drawings

FIG. 1A is a schematic diagram showing the organization of receptor-kinase fusion proteins of the
25 invention. FIG. 1B shows flow cytometry results for CD16/7/zeta, CD16/7/Lck, CD16/7/Fyn(T), CD16/7/Syk and CD16/7/ZAP-70 expressed by vaccinia recombinants in Jurkat cells. FIG. 1C shows in vitro kinase activity assay of immunoprecipitated CD16/7/zeta (negative
30 control), CD16/7/Lck, CD16/7/Fyn(T), CD16/7/Syk and CD16/7/ZAP-70; the lower molecular mass species appearing in the Fyn chimera immunoprecipitate has yet to be identified.

- 33 -

FIG. 2 shows the cytosolic calcium response triggered by crosslinking f kinase chimeras in TCR-negative Jurkat cells. The relative intracellular calcium concentration of the positive population (measured by the ratio of Indo-1 violet to blue fluorescence) is shown. Jurkat cells infected with vaccinia recombinants expressing the different fusion proteins were exposed to anti-CD16 mAb 3G8 followed by phycoerythrin conjugated goat F(ab')₂ antibodies to mouse IgG at time 0. Chimeras based on TCR zeta chain and FcRIIB2 serve as positive and negative controls respectively.

FIG. 3 shows the premature engagement of calcium response in TCR positive cells expressing Syk kinase chimera. Infection and analysis were performed as described above for Fig. 2. A substantial proportion of cells expressing Syk chimera showed a high ratio of violet to blue fluorescence prior to addition of primary antibody.

FIG. 4A and 4B show an anti-hybridoma killing assay.

FIG. 4A shows the percent ⁵¹Cr-chromate released from hybridoma target cells shown as a function of the ratio of effector cells (CTL expressing kinase chimera) to target cells; cells expressing receptor chimeras bearing the intracellular domains of TCR zeta chain and FcRIIB2 serve as positive and negative controls respectively.

FIG. 4B shows specificity of the killing (absence of bystander killing). BW5147 cells (lacking surface anti-CD16 antibody) were loaded with ⁵¹Cr-chromate and exposed to CTL expressing kinase chimeras under the same conditions as for a parallel sample of chromate-loaded 3G8 cells (expressing anti-CD16 antibody). No detectable release of chromate was observed from the BW5147 cells.

- 34 -

FIG. 5A, 5B, and 5C show that coexpression of ZAP-70 and Fyn or Lck allows induction of cytolysis and reduces latency for the calcium response. CTL were coinfectd with vaccinia recombinants expressing the indicated chimeras and analyzed for cytolytic potential or calcium mobilization. The efficacy of the chimeras is underestimated by this analysis since the fraction of cells expressing both chimeras was not independently measured (the fraction of cells expressing at least one chimera was used to normalize activity). FIG. 5A shows a cytolysis assay using CTL expressing pairs of CD16/7/kinase chimeras. FIG. 5B shows calcium response of TCR negative cells expressing pairs of CD16/7/kinase chimeras. FIG. 5C shows a cytolysis assay of CTL coexpressing a CD4/CD7/Fyn chimera and a CD16/CD7/ZAP-70 chimera. CD16/7/zeta chimera serves as the positive control, while CD16/7/FcRIIB2 chimera serves as the negative control.

FIG. 6A and 6B show that chimeras bearing kinase deletions or point mutations are ineffective in calcium mobilization and redirected cytolysis. Kinase negative fusion protein variants were constructed by deletion (in the case of Syk) or point mutation (in the case of ZAP-70) and tested for calcium response and cytolysis. FIG 6A shows calcium response in TCR negative cells. FIG. 6B shows a redirected cytolysis assay.

FIG. 7A, 7B, and 7C show that chimeras based on human Syk are essentially equipotent with chimeras based on porcine Syk. FIG. 7A is the sequence of human Syk and comparison with porcine Syk; the first 11 and last 7 residues are determined by the primer sequences. FIG. 7B shows calcium mobilization analysis of TCR negative cells expressing human Syk chimera. FIG. 7C shows a redirected cytolysis assay of CTL expressing human Syk chimera.

- 35 -

FIG. 8 shows changes in tyrosine phosphorylation pattern following crosslinking of chimeric kinases. T cell antigen receptor-negative Jurkat cells expressing the indicated chimeras or pairs of chimeras were treated with anti-CD16 and goat anti-mouse IgG second antibody and then lysed, fractionated on a polyacrylamide gel, transferred to nitrocellulose and probed with anti-phosphotyrosine antibody. Lanes marked '+' represent extracts from cells subjected to crosslinking, while those marked '-' were lysed directly without prior exposure to secondary antibody. Control lanes were created by similar treatment of TCR-negative cells expressing a CD16/7 fusion protein which did not contain an intracellular domain. For comparison, the effects of anti-CD3 treatment of TCR-positive Jurkat cells (with or without wild type vaccinia virus infection) are shown at right. The prominent bands in the vicinity of 100kD on the left part of this panel correspond to the expected molecular masses of the kinase chimeras.

FIG. 9 shows tyrosine phosphorylation of phospholipase C- γ 1 following aggregation of chimeras. PLC- γ 1 was immunoprecipitated from cells subjected to antibody crosslinking and the immunoprecipitates were fractionated on gels, transferred to nitrocellulose, and probed with anti-phosphotyrosine antibody. A substantial increase in phosphorylated PLC- γ 1 was seen following aggregation of Syk chimeras, whereas a more limited but easily detectable increase is seen following coaggregation of Fyn and ZAP-70 chimeras.

FIG. 10A and 10B show in vitro kinase assays. Cells expressing chimeric kinases were subjected to antibody-mediated chimera crosslinking, after which the kinases were immunoprecipitated and the immunoprecipitates evaluated for phosphorylation of endogenous substrate. FIG. 10A shows a comparison of the

- 36 -

activity of immunoprecipitated kinase chimeras over an incubation period of ten minutes, using immunoprecipitates isolated from crosslinked (+) or uncrosslinked (-) cells. FIG. 10B shows a time course of
5 assimilation of phosphate label into endogenous substrates by Syk kinase chimera, with (+) or without (-) crosslinking.

T Cell Activation by Clustered Tyrosine Kinases

There now follows a description of particular
10 embodiments of the invention. In this description, it is demonstrated that nonreceptor kinases are activated by simple clustering events. Artificial receptor kinases were created whose intracellular domains consisted of the complete Src or Syk family kinase sequences and examined
15 for the consequences of aggregation by external crosslinking stimuli. A clear distinction emerged between the Syk and Src family kinase activities: crosslinking the latter did not lead to significant cellular activation, while crosslinking the former led to
20 the appearance of free intracellular calcium ion and, in the case of Syk, of cytolytic potential. The failure of ZAP-70 chimeras to induce distal receptor mediated programs could be overcome by coclustering ZAP-70 chimera with either Fyn or Lck kinase chimeras. The examples now
25 described are provided for the purpose of illustrating, not limiting, the invention.

Construction of Protein-Tyrosine Kinase Chimeras and Demonstration of Efficacy

Gene fusions encoding proteins resembling cell
30 surface receptor kinases were constructed by appending a DNA fragment encoding the extracellular domain of the

- 37 -

CD16 molecule to a short spacer segment encoding the juxtamembraneous and transmembrane domains of CD7 joined in turn to the complete coding sequences of the human Lck (Koga et al., 1986, Eur. J. Immunol. 16:1643-1646),
5 murine Fyn (T) (Cooke and Perlmutter, 1989, New. Biol. 1:66-74), porcine Syk (Taniguchi et al., 1991, J. Biol. Chem. 266:15790-15796) and human ZAP-70 (Chan et al., 1992, Cell 71:649-662) tyrosine kinases (Fig. 1A). The resulting tripartite gene fusions were introduced into
10 recombinant vaccinia viruses by homologous recombination and selection for coexpression of the E. coli gpt gene product. Infection of cells with the recombinants resulted in the efficient cell surface expression of all four kinase chimeras (Fig. 1B). Immunoprecipitation of
15 the resulting protein chimeras with anti-CD16 antibodies revealed the presence of molecular species of the expected masses which were active in an in vitro phosphorylation assay (Fig. 1C).

We next examined whether crosslinking of the
20 fusion proteins would allow the accumulation of free intracellular calcium in a fashion similar to that found with fusion proteins based on T cell antigen receptor intracellular domains. To do this we infected various cells with vaccinia recombinants and measured the
25 relative cytoplasmic calcium concentration following crosslinking of the extracellular domains with antibodies. Both spectrofluorimetric (bulk population) and flow cytometric (single cell) measurements were performed, with cells loaded with the dye Indo-1
30 (Gryniewicz et al., 1985, J. Biol. Chem. 260:3440-3450; Rabinovitch et al., 1986, J. Immunol. 137:952-961). Flow cytometric analyses were performed on data obtained from cells whose cell surface expression of CD16, as determined by phycoerythrin fluorescence intensity, fell
35 within a relatively narrow predefined range. Although

- 38 -

minor variations in mean fluorescence intensity were still observed within this range (due to differences in the underlying distribution of chimeras expressed by the cells), this approach allowed us to contrast the responses of cells bearing approximately the same number of receptors. Figure 2 shows an analysis of data collected from cells of a mutational variant of the Jurkat human T cell leukemia line lacking T cell antigen receptor (Weiss and Stobo, 1984, J. Exp. Med. 160:1284-1299). In these cells neither Lck nor Fyn chimeras had the capacity to mobilize calcium following crosslinking. In several experiments clustering of the Lck fusion protein resulted in a slight decrease in resting calcium concentration, relative to the negative control, a fusion protein based on the low affinity IgG receptor FcRIIB2 intracellular domain (Kolanus et al., 1992, EMBO J. 11:4861-4868). Aggregation of fusion proteins based on both ZAP-70 and Syk was highly effective in promoting the appearance of free cytoplasmic calcium ion, roughly as effective as aggregation of a similar chimera bearing the intracellular domain of the T cell receptor zeta chain. A slight delay in onset of the calcium response was seen with both ZAP-70 and Syk kinase chimeras, relative to the time of onset of calcium mobilization by zeta chimera. In T cell receptor positive cells (Fig. 3), flux evaluation of Syk chimeras was partially confounded by a high resting concentration of free calcium ion, suggestive of a constitutive engagement of the calcium regulatory apparatus.

Introduction of the chimeras into a cytolytic T cell line then allowed us to assess the fusion proteins' potential to engage effector function. In this assay, anti-CD16 hybridoma cells which express cell surface IgG antibody against CD16 (Fleit et al., 1982, Proc. Natl. Acad. Sci. USA 79:3275-3279; Shen et al., 1989, Mol.

- 39 -

Immunol. 26:959-969) are used as target cells. The hybridoma cells are labeled by incorporation of ^{51}Cr -chromate and cosedimented with effector cells, prepared by infection of a human allospecific cytotoxic T lymphocyte (CTL) line with vaccinia recombinants expressing the CD16/CD7/kinase fusion proteins. Expression of the chimeric receptor kinases allows the infected CTL cells to bind to the target cells, and if competent, to lyse them, a process which is measured by release of the incorporated ^{51}Cr -chromate. The relative potency of the armed CTL is determined by comparison of the ratio of effector to target cells needed to achieve a given proportion of ^{51}Cr release. Fig. 4A shows that CTL expressing chimeric receptors comprising the Src family kinases Lck or Fyn (T), or the Syk family kinase ZAP-70, are incapable of mediating cytolysis against anti-CD16 hybridoma targets. However CTL expressing a kinase chimera based on the Syk protein product were essentially as effective as CTL expressing a chimera composed of CD16 fused in the same manner to the intracellular domain of the T cell receptor zeta chain. The cytolytic activity directed by the CD16/CD7/kinase chimeras could not be ascribed to a nonspecific release of cytotoxic granules because cosedimentation of an irrelevant chromium-laden target with the kinase-armed CTL did not result in detectable release of labeled chromium (Fig. 4B).

The disparity between Syk and ZAP-70 activities in the cytolysis assay was unexpected in light of the similar activities of the two chimeras in the calcium response assay. In light of the demonstration that coexpression of nonchimeric ZAP-70 and Src-family kinases led to activation in COS cells (Chan et al., 1992, Cell 71:649-662), we undertook an evaluation of the relative potential of pairs of kinase chimeras to effect cytolysis. CTL effectors were coinfectd with

- 40 -

recombinant vaccinia viruses encoding ZAP-70 and Lck chimeras, ZAP-70 and Fyn (T) chimeras, or Lck and Fyn (T) chimeras. Fig. 5A shows that the coexpression of ZAP-70 and Fyn (T) chimeras, or ZAP-70 and Lck chimeras, endowed CTL with an activity essentially equipotent with that of CTL expressing CD16/CD7/Syk kinase chimeras alone, an activity in turn as potent as that displayed by CD16/CD7/zeta chimeras. Coexpression of Lck and Fyn(T) chimeras did not allow significant cytolytic potential to be redirected against anti-CD16 target cells (Fig. 5A). Evaluation of the calcium mobilization potential of cells coinfecting with pairs of kinase fusion proteins showed that coexpression of ZAP-70 and Src family kinase chimeras increased the rapidity with which calcium was mobilized in response to receptor crosslinking and that coexpression of Lck and Fyn(T) chimeras did not result in a significant accumulation of free intracellular calcium (Fig. 5B). To further explore the role of the Fyn chimera in the activation response induced by coaggregation, we prepared a Fyn chimera consisting of the extracellular and transmembrane domains of CD4 fused to Fyn in a manner similar to that of the CD16 chimeras. Fig. 5C shows that the effectiveness of this chimera in the directed cytolysis assay is ten to twenty fold lower than that of the comparable CD16 chimera, suggesting that physical association of the chimeric kinases is important for activation. However the cytolytic activity of cells expressing the two chimeras is significantly greater than would be observed for cells expressing ZAP-70 chimera alone. Because of the relatively high level of kinase expression in this system, it cannot be excluded that the residual activity reflects spontaneous random association of the CD4/Fyn chimera with CD16/ZAP-70.

- 41 -

To establish that the activation seen in both calcium response and cytolysis assays was directly attributable to the relevant kinase activity and not to passive association of the kinases with existing signal transduction elements whose indirect aggregation then initiated the activation response, we created kinase negative variants of both the porcine Syk and human ZAP-70 receptor chimeras. Fig. 6 shows that receptor chimeras lacking either substantially all of the kinase domain (in the case of Syk) or bearing a point mutation abrogating phosphotransferase activity (in the case of ZAP-70) lacked in vitro kinase activity and were incapable of mediating calcium mobilization following crosslinking, or of mediating receptor redirected cytolysis.

Because the interaction of porcine Syk with the human cellular apparatus might not be identical to the interaction of human Syk, we also constructed similar protein chimeras based on human Syk, after isolating human Syk sequences by PCR with primers corresponding to the amino and carboxyl termini of the porcine protein sequence. Fig. 7A shows that pig and human Syk are strikingly similar proteins, as suggested by analysis of PCR products corresponding to portions of the kinase and second SH2 domains (Chan et al., 1992, Cell 71:649-662). Consonant with this, human Syk chimeric receptor proteins behaved essentially identically to the porcine constructs in calcium release and cytolysis assays (Fig. 7B and 7C).

To establish whether aggregation of the chimeric tyrosine kinases results in a significant change in the abundance of phosphotyrosine proteins, T cell receptor negative cells were infected with vaccinia recombinants encoding the chimeric kinases. The extracellular domains of the chimeras were crosslinked with antibodies, and total cellular lysates of the activated cells were

- 42 -

fractionated by electrophoresis, transferred to membranes and analyzed with an antibody recognizing phosphotyrosine. Lysates were prepared by disruption of cells in nonionic detergents in the presence of vanadate
5 with or without EDTA, or by lysis in the presence of sodium dodecyl sulfate, followed by sonication to shear the liberated DNA. The pattern of phosphotyrosine proteins was different in each case and lysates prepared with vanadate but without EDTA showed additional species
10 not present in lysates prepared in SDS alone, suggesting that EDTA may inhibit the postlysis action of tyrosine kinases as well as phosphatases. The use of direct lysis in SDS was found to lead to more reproducible patterns of protein tyrosine phosphorylation than lysis with nonionic
15 detergents in the presence of EDTA and vanadate. Fig. 8 shows that aggregation of chimeras bearing Syk, ZAP-70, or Fyn plus ZAP-70 results in the appearance or increased phosphorylation of several protein species comigrating with proteins which show increased phosphorylation
20 following antigen receptor crosslinking. In particular, the pattern of bands induced by Syk chimera clustering is very similar to the pattern induced by anti-CD3 antibody in T cell receptor-positive cells (Fig. 8). Among these is an approximately 150 kD protein induced by aggregation
25 of Syk chimera, but also induced by coaggregation of Fyn and ZAP-70 chimeras. In preliminary experiments this phosphoprotein was observed to comigrate with phospholipase C- γ (data not shown).

To establish the effects of kinase chimera
30 clustering on PLC- γ directly, we crosslinked chimeras, precipitated PLC- γ with a mixture of monoclonal antibodies, and analyzed the resulting immunoprecipitates for the presence of phosphotyrosyl proteins. Fig. 9 shows that clustering of Syk results in a substantial
35 increase in the tyrosine phosphate content of PLC- γ ,

- 43 -

while cocrosslinking of Fyn plus ZAP-70 chimeras results in a less dramatic but easily detectable increase in tyrosine phosphate. Cells expressing Fyn chimera alone showed a modest basal phosphorylation of PLC- γ which did not increase following receptor aggregation (Fig. 9).

Whether the same tyrosine residues are phosphorylated by Syk, Fyn, or ZAP-70 plus Fyn, is presently unknown. Because cells expressing Fyn chimera showed neither resting nor induced calcium mobilization, the phosphotyrosine signal seen in these cells may represent utilization of other sites on PLC- γ than those which mediate phospholipase activation.

In a preliminary attempt to account for the changes in phosphotyrosine pattern we evaluated the activity of the various kinases following clustering in an in vitro autophosphorylation assay. Chimeras were aggregated, immunoprecipitated, and evaluated for their ability to incorporate phosphate label into the protein species present in the immunoprecipitate. Fig. 10A shows that under these conditions no increase in kinase activity was detected following crosslinking when the kinase assay was carried out for ten minutes. Although incorporation of labeled phosphate continues to increase for up to 30 minutes in this assay, it is unclear what factors limit activity; in particular the observed kinetics may be dominated by the rate of dissociation of the immune complexes, allowing kinase diffusion to unutilized substrate. In an attempt to control for this effect, as well as to maximize the sensitivity of the assay, we also evaluated Syk kinase activity with or without prior crosslinking over a very brief time course from 5 to 30 seconds; however here too there was no significant increase in kinase activity (Fig. 10B). Although at present we cannot exclude the possibility that an increase in activity would be demonstrable with

- 44 -

an appropriate substrate, an aggregation-induced increase in Syk chimera activity was also not observed when an exogenous peptide substrate (a Y 19 K substitution of cdc 2 residues 6-20) was used to measure kinase activity.

5 Possible Mechanisms

The mechanism by which a simple physical stimulus, receptor aggregation, results in the transmission of a distinctive chemical signal to immune system cells remains unknown. Previous studies have established that

10 Src-family kinases can be found associated with many of the important aggregation-activated receptors of the immune system, and that crosslinking of such receptors frequently leads to increased kinase activity. More recently, the related Syk and ZAP-70 kinases have been

15 found to associate either stably (in the case of Syk) or transiently (in the case of ZAP-70) with the B and T cell antigen receptors, respectively, and at least in the case of Syk, receptor crosslinking has been reported to result in increased kinase activity.

20 In this work we have shown that aggregation of Syk family kinases, but not the Src family kinases Lck or Fyn, leads to a calcium response in cells lacking T cell receptor. The response appears not to be due to an indirect association with other T cell receptor or signal

25 transduction components because kinase negative mutants can not induce the calcium response. Aggregation of chimeras containing the Syk kinase is sufficient to allow induction of specific cytolysis, while induction of similar cytolysis by ZAP-70 chimera requires the

30 additional participation of a Src family kinase. At present it is unclear which of the Syk family kinases is likely to play a more important role in T cell activation: both ZAP-70 and Syk are expressed in T cells,

- 45 -

including the cell lines used in this study, and at least on responsive human T cell line contains Syk but not ZAP-70 as judged by specific PCR amplification products. The pattern of increased protein tyrosine phosphorylation
5 seen following Syk chimera clustering is very similar to the pattern seen after crosslinking of the T cell antigen receptor.

One simple model for activation of immune system cells by nonreceptor tyrosine kinases invokes a
10 receptor-associated kinase whose aggregation leads to activation either by physical association (e.g., by forming active kinase dimers) or by mutual enzymatic action (e.g., crossphosphorylation); the activated enzyme then acts on key intracellular substrates required for
15 calcium mobilization and inositolphosphate synthesis. Support for this sequence of events can be found in studies reporting increases in receptor-associated kinase activity following receptor crosslinking (e.g., Bolen et al., 1991, Adv. Cancer Res. 57:103-149; Burkhardt et al.,
20 1991, Proc. Natl. Acad. Sci. USA 88:7410-7414; Eiseman and Bolen, 1992, Nature 355:78-80; Hutchcroft et al., 1992, Proc. Natl. Acad. Sci. USA 89:9107-91111/J. Biol. Chem. 267:8613-8619; Tsygankov et al., 1992, J. Biol. Chem. 267:18259-18262; Wong et al., 1992, Oncogene
25 7:2407-2415). However the reported changes in kinase activity are in most cases modest, and contrast with the dramatic changes in phosphotyrosyl protein pattern seen in vivo. Because it is difficult to unambiguously rule out activation by weak allosteric interactions using in
30 vitro kinase assays as a tool, we cannot at this point make a definitive statement about the relative importance of kinase activation in the initiation of signal transduction. But the data presented here suggest that aggregation-induced repartitioning of enzyme and
35 substrate may be an important factor in directing an

- 46 -

existing activity toward the appropriate physiological target.

Aggregation of chimeras based on Syk family kinases led to a calcium response which was slightly
5 delayed but similar in amplitude to the response seen following aggregation of zeta receptor chimeras in cells lacking endogenous T cell receptor. A more profound delay in the appearance of free calcium ion was observed following ZAP-70 chimera crosslinking, and this delay
10 could be substantially abolished by cocrosslinking ZAP-70 and Fyn chimeras. At present the explanation for the observed latency is unclear. Because cocrosslinking accelerated calcium mobilization, it is tempting to ascribe the delay to the relative inefficacy of ZAP-70
15 for aggregation-mediated autoactivation. But other factors may be equally important, and the tethering of ZAP and Syk kinases at the cell surface may actually be an impediment to activation when compared to the normal process; for example if in the normal course of events
20 Syk family kinases are transiently recruited to clustered receptors, activated and then released to diffuse to their substrates, the permanent linkage of the kinase domain to the plasma membrane could be restrictive by hindering access to substrate and/or limiting signal
25 amplification due to the inability of the chimeric receptors to function as a kind of catalytic center for kinase activation.

A second peculiarity of the calcium response was the finding that T cell receptor positive cells
30 expressing chimeras based on human or porcine Syk showed a high baseline concentration of free calcium ion, suggesting that calcium release had been spontaneously triggered. A similar finding was not observed in receptor negative cells. This result runs counter to a
35 general trend we have observed, in which T cell receptor

- 47 -

negative mutants of human T cell tumor lines typically are hyperresponsive to exogenously introduced trigger molecules. To account for the apparent requirement for T cell receptor in the spontaneous activation process, we
5 propose two possible and related explanations. One is that chimeric Syk kinase may act constitutively on T cell receptor intracellular domains to create phosphotyrosine targets for the SH2 domains of the receptor chimera, leading to intracellular aggregation through a
10 multivalent T cell receptor bridge, in turn promoting kinase activation. Another possibility is that the T cell receptor negative cell line may have lower levels of a membrane associated kinase required for activation of Syk, either because a global regulatory circuit results
15 in decreased de novo synthesis of the hypothetical kinase, or because the absence of antigen receptor results in a disregulated intracellular trafficking.

In B cells, the Syk kinase has been reported to be constitutively associated with the intracellular elements
20 of the IgM antigen receptor (Hutchcroft et al., 1992, Proc. Natl. Acad. Sci. USA 89:9107-91111; Hutchcroft et al., 1992, J. Biol. Chem. 267:8613-8619). The exact mechanism of this association is unclear, but one possibility is that phosphotyrosine is not required for
25 interaction of Syk SH2 domains with the tyrosine trigger motif present in the cytoplasmic domain of the mb-1 and B 29 receptor chains. A partial precedent for this suggestion is the report that the Philadelphia chromosome breakpoint cluster region gene product BCR binds to a
30 variety of SH2 domains in a phosphotyrosine independent manner (Pendergast et al., 1991, Cell 66:161-171; Muller et al., 1992, Mol. Cell. Biol. 12:5087-5093). In this case, though, it appears likely that phosphoserine and/or phosphothreonine residues play a critical role in the
35 interaction. Alternatively, Syk may associate with IgM

- 48 -

receptor intracellular motifs through the unique region located between the Syk SH2 elements and the catalytic domain. A third possibility is that only a very small amount of tyrosine phosphorylated peptide is necessary to
5 recruit functionally important levels of Syk to the inner face of the plasma membrane, and this low level of tyrosine phosphate has thus far escaped detection.

Although in B cells the requirement for a Src family kinase in activation has not been definitively
10 established, in T cells two kinases, Lck and Fyn (T) have been shown by somatic or organismic genetics to play important roles (Appleby et al., 1992, Cell 70:751-763; Karnitz et al., 1992, Mol. Cell. Biol. 12:4521-4530; Stein et al., 1992, Cell 70:741-750; Straus and Weiss,
15 1992, Cell 70:585-593). At present we cannot conclusively establish whether the action of these kinases normally precedes or follows the action of the Syk family kinases. One hypothesis accounting for the action of ZAP-70 in T cell activation invokes a
20 receptor-associated Src family kinase whose aggregation permits a transitory phosphorylation of receptor chains in turn leading to association of ZAP-70 and subsequent cellular activation. The initial phosphorylation of receptor chains proposed in this model must be
25 distinguished from the stable phosphorylation of zeta seen at longer times following receptor crosslinking.

In murine T cells a small proportion of T cell receptor complexes contain a zeta-related molecule called eta (Baniyash et al., 1988, J. Biol. Chem. 263:9874-9878)
30 which represents an alternatively spliced form (Clayton et al., 1991, Proc. Natl. Acad. Sci. USA 88:5202-5206). Eta differs from zeta at the carboxyl terminus, and lacks the most distal of the six tyrosines found in murine zeta (Jin et al., 1990, Proc. Natl. Acad. Sci. USA 87:3319-
35 3323). Although the phosphorylation of the zeta chain of

- 49 -

murine TCR zeta-zeta isoforms can readily be detected following antibody-mediated receptor crosslinking, under similar circumstances the TCR eta chain is not detectably phosphorylated (Bauer et al., 1991, Proc. Natl. Acad. Sci. USA 88:3842-3846). Stable phosphorylation of the zeta chain appears to require two closely apposed zeta chains since TCR isoforms bearing zeta-eta heterodimers are not phosphorylated following receptor crosslinking (Bauer et al., 1991, Proc. Natl. Acad. Sci. USA 88:3842-3846). Despite the differences in phosphorylation, cell lines comprising TCR isoforms consisting solely of eta homodimers are functionally indistinguishable from cell lines bearing only zeta homodimers (Bauer et al., 1991, Proc. Natl. Acad. Sci. USA 88:3842-3846). Thus phosphorylation of zeta, as observed 30 minutes after antibody mediated receptor aggregation, does not correlate with activation. In a separate study, examination of the time course of accumulation of phosphotyrosine phosphoproteins following TCR aggregation has shown that the earliest observable species are two proteins of mass 135 and 100 kD, whose phosphorylation is first detectable at five and fifteen seconds after crosslinking, respectively, and whose half maximal phosphorylation, in both cases at approximately 30 seconds, precedes the times of half maximal calcium mobilization and inositol phosphate formation (June et al., 1990, Proc. Natl. Acad. Sci. USA 87:7722-7726; June et al., 1990, J. Immunol. 144:1591-1599). By contrast, the rate of phosphorylation of zeta is substantially slower, leading to half maximal substitution at approximately three to five minutes post-stimulation, well after calcium accumulation and liberation of inositol phosphates are observed (June et al., 1990, Proc. Natl. Acad. Sci. USA 87:7722-7726; June et al., 1990, J. Immunol. 144:1591-1599).

- 50 -

Thus if the two step model is correct, the tyrosine phosphorylation necessary to recruit ZAP-70 to the inner face of the plasma membrane must be a more rapid, presumably transitory, event than observed in the studies above. Recently it has been suggested that tyrosine phosphorylation with an appropriately fast (ca. fifteen second) onset can be detected on both zeta and CD3 epsilon chains following receptor crosslinking (Wange et al., 1992, J. Biol. Chem. 267:11685-11688), and that a 70 kD protein bearing tyrosine phosphate can be found associated with both zeta and epsilon chains. It is not presently clear whether a stable association with phosphorylated receptor chains is a prerequisite for successful T cell activation.

As a general assertion, though, the results reported here suggest that Syk family kinases act more directly on the effector apparatus of T cells than Src family kinases. An increasing body of evidence suggests that Src family kinases can associate with a number of cell surface molecules which are not members of the antigen/Fc receptor family, including CD2 (Bell et al., 1992, Mol. Cell. Biol. 12:5548-5554), CD23 (Sugie et al., 1991, Proc. Natl. Acad. Sci. USA 88:9132-9135), CD36 (Huang et al., 1992, J. Biol. Chem. 267:5467-5473), IL-2 receptor beta chain (Hatakeyama et al., 1991, Science 252:1523-1528) and various phosphatidylinositol anchored proteins (Stefanova et al., 1991, Science 254:1016-1019; Thomas and Samelson, 1992, J. Biol. Chem. 267:12317-12322), some of which are known to require the additional presence of the antigen receptor to promote activation in T cells. A simple explanation for the latter requirement maybe that the trigger motifs on the antigen receptor act as substrates for Src family kinases, allowing the subsequent docking of Syk family kinases, followed perhaps by some modifying event promoting their

- 51 -

activation. Given the difficulty of establishing a causal chain of phosphorylation and activation it may also be that the trigger motifs have only a transitory role, to act as a kind of catalytic center for the
5 recruitment, activation, and release of effector kinases.

Src family kinases are broadly distributed throughout nonhematopoietic lineages, and recent studies using Fyn negative mice have shown a role for Fyn in the sustenance of long term potentiation, the phenomenon of
10 facilitated synaptic transmission thought to underlie the initial consolidation of associative memory (Grant et al., 1992, Science 258:1903-1910). If similar activation pathways are mediated by Src family kinases in other cell types, Syk family kinases may also prove to be more
15 extensively distributed throughout the extrahematopoietic compartments.

Experimental Methods

Construction of chimeras. The entire coding regions of the human Lck (Koga et al., 1986, Eur. J. Immunol.
20 16:1643-1646), murine Fyn (T) (Cooke and Perlmutter, 1989, New. Biol. 1:66-74), porcine Syk (Taniguchi et al., 1991, J. Biol. Chem. 266:15790-15796) and human ZAP-70 (Chan et al., 1992, Cell 71:649-662) kinases were attached to the intracellular domain of a chimeric
25 transmembrane protein consisting of the CD16 extracellular domain joined to a short 'stalk' segment and transmembrane domain of CD7. The CD7 intracellular domain was truncated at the stop transfer sequence by addition of an Mlu site. The various kinases were
30 adapted with an Mlu site in the appropriate reading frame to allow a tripartite fusion protein to be expressed. The pig Syk sequence was obtained by reverse transcription and PCR of total pig lymphocyte RNA using

- 52 -

primers bearing appropriate restriction sites. ZAP-70 sequences were similarly obtained by PCR from a human T cell cDNA library. Several isolates were sequenced in parallel and a mutation-free coding sequence was derived for each kinase by restriction fragment interchange. The resulting coding sequences were inserted into a vaccinia virus expression vector downstream from the CD16/CD7 sequences. Human Syk was isolated from a natural killer cell cDNA library and from a Daudi cell library using primers corresponding to the ends of the porcine sequence. The forward primer bore the sequence atg gca gac agt gcc aac cac ttg ccc ttc ttc t and the reverse primer bore the sequence cgc ggg gcg gcc gct tta att cac cac gtc gta gta gta. After initial amplification revealed the presence of bands of the expected size a reamplification (10 cycles) was performed using an extension primer at the 5' end having the sequence cgc ggg acg cgt acc atg gca gac agt gcc aac, allowing the fragment to be ligated to an Mlu I-cut vector.

Calcium mobilization analysis. Flow cytometric and bulk spectrophotometric analyses were conducted on cells expressing recombinant kinases using the calcium sensitive fluorophore Indo-1 as previously described (Romeo and Seed, 1991, Cell 64:1037-1046; Romeo et al., 1992, Cell 68:889-897). Briefly, cells of the Jurkat mutant subline JRT 3.T 3.5 (Weiss and Stobo, 1984, J. Exp. Med. 160:1284-1299) were infected with recombinant vaccinia viruses for one hour in serum free IMDM at an moi of 10 and incubated for three to nine hours in IMDM, 10% FBS. Cells were collected by centrifugation and resuspended at 3×10^6 /ml in complete medium containing 1mM Indo-1 acetomethoxyester (Grynkiewicz et al., 1985, J. Biol. Chem. 260:3440-3450) (Molecular Probes, Eugene, OR) and incubated at 37°C for 45 minutes. The Indo-1 loaded cells were pelleted and resuspended at 1×10^6 /ml

- 53 -

in serum free IMDM and stored at room temperature in the dark. Cells were analyzed for free calcium ion by simultaneous measurement of the violet and blue fluorescence emission by flow cytometry (Rabinovitch et al., 1986, J. Immunol. 137:952-961). To initiate calcium flux, either unconjugated 3G8 (anti-CD16) monoclonal antibody (at 1 μ g/ml) was added to the cell suspension followed by 10 μ g/ml of phycoerythrin (PE)-conjugated Fab 02 goat anti-mouse IgG at time 0, or a PE-conjugated anti-CD4 antibody (Leu-3a, Becton Dickinson) was added, followed by unconjugated second antibody. Histograms of the violet/blue emission ratio were collected from the PE positive (infected) cell population, which typically represented 40-80% of the cells. The violet/blue emission ratio prior to the addition of antibody was used to establish the normalized initial ratio, set equal to unity.

Lymphocyte cytotoxicity assay. A CD8⁺ CD4⁻ HLA B44 restricted cytolytic line (WH3) was maintained in IMDM, 10% human serum with 100 U/ml of IL-2 and was periodically stimulated with irradiated (3000 rad) mononuclear cells having the HLA B44 haplotype. Cells were grown for at least 10 days following stimulation before use in cytotoxicity assays. The cells were infected with recombinant vaccinia at a multiplicity of infection of at least 10 for one hour in serum free medium, followed by incubation in complete medium for three hours. Cells were harvested by centrifugation and resuspended at a density of 1×10^7 /ml. 100 μ l were added to each well of a U-bottom microtiter plate containing 100 μ l/well of complete medium. Cells were diluted in two-fold serial steps. Two wells for each sample did not contain lymphocytes, to allow spontaneous chromium release and total chromium uptake to be measured. An aliquot of 10^6 3G8 10-2 target cells (Shen

- 54 -

et al., 1989, Mol. Immunol. 26:959-969) or was centrifuged and resuspended in 50 μ l of sterile ^{51}Cr sodium chromate (1 m Ci/ml, DuPont) for one hour at 37°C with intermittent mixing, then washed three times with 5 PBS. 100 μ l of labelled cells resuspended in medium at 10^5 /ml were added to each well. The microtiter plate was spun at 750 x g for 1 minute and incubated for 4 hours at 37°C. At the end of the incubation period the cells in each well were resuspended by gentle pipetting, a sample 10 removed to determine the total counts incorporated, and the microtiter plate was spun at 750 x g for 1 minute. 100 μ l aliquots of supernatant were removed and counted in a gamma rays cintillation counter. The effector to target ratio was corrected for the percent of effector 15 cells infected (usually >70%).

Creation of mutant kinase chimeras. A porcine Syk kinase negative fusion protein variant was created by cleavage of the chimera with Stu I and Not I (the latter lying just 3' to the carboxyl terminus), filling in the 20 Not I site, and ligating the ends together. The resulting sequence joined the first 298 residues of pig Syk to 4 extraneous residues (GPRL) before terminating. A point mutation (K369G) in the ATP binding site of ZAP-70 was created by insertion of a duplex 25 oligonucleotide fragment between the BallI and EarI sites located between nucleotide residues 1319 and 1345 of the top strand of the sequence reported by Chan et al. (1992, Cell 71:649-662). The resulting sequence encoded glycine at residue 369 in place of lysine.

30 Immunoprecipitation and kinase assay. Approximately 2×10^6 Hela S3 cells were infected for one hour in serum free DME medium with recombinant vaccinia at a multiplicity of infection of at least ten. 5 hrs after infection the cells were harvested, washed twice with 35 phosphate buffered saline and lysed in 1% Triton X-100,

- 55 -

0.15 M NaCl, 0.02 M HEPES pH 7.3, 5 mM EDTA, 5 mM NaF, 0.2 M MNaVO_3 , 10 $\mu\text{g/ml}$ leu peptin, 10 $\mu\text{g/ml}$ a protinin and 1mM PMSF. After a 10 min. incubation on ice, the nuclei were removed by centrifugation and the CD16 fusion proteins immunoprecipitated with antibody BMA209/2 and protein-A sepharose. The fusion protein loaded resin was washed 3 times with lysis buffer followed by a final wash with 20 mM Hepes pH 7.3. To each sample was added 10 μl of kinase buffer (20 mM Hepes pH 7.3, 10 mM MgCl_2 , 10 mM MnCl_2) containing 10 μCi of $[\gamma\text{-}^{32}\text{P}]\text{ATP}$ ($>3000 \text{ Ci/mmol}$). The reactions were allowed to incubate at room temperature for 10 min. and terminated by the addition of 20 μl of 2X sample loading buffer (4% SDS, 100 mM Tris pH6.8, 20% glycerol, and 10% β -mercaptoethanol). After the samples were boiled for 3 min., aliquots were run on a 4-15% gradient gel. Kinase assays with a soluble peptide substrate corresponding to positions 6-20 of cdc 2, in which tyr 20 was replaced with lys, were performed according to the manufacturer's recommendations (UBI).

Immunoblot analysis of protein tyrosine phosphorylation. TCR negative 3.5 cells were infected with recombinant virus stocks (moi of at least 10) for one hour. Cells were subsequently incubated at 37°C for 8-12h, centrifuged, washed and resuspended in Iscove's medium without serum at 10^7 cells per ml. Aliquots of cells were incubated with anti-CD16 mAb (3G8, Medarex or BMA209/2, Behringwerke) at 1 μg antibody per $2 - 3 \times 10^6$ cells. Stimulated samples were further incubated with a 3-5 fold excess of an affinity purified anti-mouse IgG 1 antibody (Southern Biotechnology) for 5 min. Cells were subsequently processed according to Secrist, J.P., Burns, L.A., Karnitz, L., Koretzky, G.A., and Abraham, R.T. (J. Biol. Chem. 268,5886-5893, 1993). with slight modifications. Incubations were terminated by adding SDS to a final concentration of 1% and samples were boiled.

- 56 -

for three minutes. DNA was sheared by sonication for 1 min using a Heat Systems Ultrasonics, Inc., 2X sample buffer was added and aliquots corresponding to 10^5 to 2.5×10^5 cells were separated on polyacrylamide gels and the proteins transferred by semidry electroblotting (Hoefer) onto nitrocellulose (Schleicher and Schuell BA45). Filters were blocked for one hour in Tris Buffered Saline with 0.05% Tween-20 (TBST) containing 1.5% chicken ovalbumin (Sigma), washed in TBST, and transferred into solution containing anti-phosphotyrosine antibody 4G10 (UBI) at a 1:10000 dilution and incubated at 22° for 1-2h. After TBST washes, filters were incubated in TBST and a 1:5000 dilution of anti-mouse horseradish peroxidase conjugate for 1 h. Phosphorylated protein bands were detected by chemiluminescence (ECL, Amersham). Exposure times varied between 2s and 5 min.

Construction of Human IgG1:Protein-Tyrosine Kinase Chimeras

Chimeric molecules may be produced which include the extracellular domain of an antibody molecule specific for a target cell and the intracellular domain of a protein-tyrosine kinase (e.g., those kinases described herein). To produce such a molecule, human IgG1 heavy chain sequences are prepared by joining sequences in the C_H3 domain to a cDNA fragment derived from the 3' end of the transmembrane form of the antibody mRNA. The 3' end fragment is obtained by polymerase chain reaction using a tonsil cDNA library as substrate, and oligonucleotides having the sequences:

CGC GGG GTG ACC GTG CCC TCC AGC AGC TTG GGC and
CGC GGG GAT CCG TCG TCC AGA GCC CGT CCA GCT CCC
CGT CCT GGG CCT CA,
corresponding to the 5' and 3' ends of the desired DNA fragments respectively. The 5' oligo is complementary to

- 57 -

a site in the C_H1 domain of human-IgG1, and the 3' oligo is complementary to a site just 5' of the sequences encoding the membrane spanning domain. The PCR product is digested with BstXI and BamHI and ligated between
5 BstXI and BamHI sites of a semisynthetic IgG1 antibody gene bearing variable and constant regions. Following the insertion of the BstXI to BamHI fragment, the amplified portions of the construct are replaced up to the SmaI site in C_H3 by restriction fragment interchange,
10 so that only the portion between the SmaI site and the 3' oligo is derived from the PCR reaction.

To create a human IgG1 chimeric receptor, the heavy chain gene ending in a BamHI site is joined to the kinase intracellular domain of interest by standard
15 techniques. Levels of chimeric receptor expression may be determined by flow cytometry. Increases in expression may be accomplished by coexpression of a plasmid encoding an antibody light chain cDNA.

To create a single transcription unit which would
20 allow both heavy and light chains to be expressed from a single promoter, a plasmid encoding a bicistronic mRNA is created from heavy and light chain coding sequences, and the 5' untranslated portion of the mRNA encoding the 78kD glucose regulated protein, otherwise known as grp78, or
25 BiP. grp78 sequences are obtained by PCR of human genomic DNA using primers having the sequences:

CGC GGG CGG CCG CGA CGC CGG CCA AGA CAG CAC and

CGC GTT GAC GAG CAG CCA GTT GGG CAG CAG

at the 5' and 3' ends respectively. Polymerase chain
30 reactions with these oligos are performed in the presence of 10% dimethyl sulfoxide. The fragment obtained by PCR is digested with NotI and HincII and inserted between NotI and HpaI sites downstream from human IgG1 coding sequences. Sequences encoding a human IgG kappa light
35 chain cDNA are then inserted downstream from the grp78

- 58 -

leader, using the HincII site and another site in the vector. The expression plasmid resulting from these manipulations consists of the semisynthetic heavy chain gene, followed by the grp78 leader sequences, followed by
5 the kappa light chain cDNA sequences, followed by polyadenylation signals derived from an SV40 DNA fragment. We have previously demonstrated that transfection of COS cells with this expression plasmid gave markedly improved expression of heavy chain
10 determinants, compared to transfection of plasmid encoding heavy chain determinants alone.

To create a bicistronic gene comprising a heavy chain/receptor chimera and a light chain, the upstream heavy chain sequences can be replaced by any chimeric
15 heavy chain/ receptor gene described herein.

Once constructed, the IgG-tyrosine kinase chimeras may be cloned into an expression vector, introduced into a host cell, and tested by any of the assays described herein (e.g., by calcium mobilization or cytolysis
20 assays).

Construction of CD4-Tyrosine Kinase Chimeras

Chimeric molecules may be produced which include the extracellular domain of the CD4 molecule and the intracellular domain of a protein-tyrosine kinase (e.g.,
25 those kinases described herein). To produce such a molecule, the tyrosine kinase-encoding sequence (for example, cDNA) is isolated (for example, as described above). This sequence is then joined to the extracellular domain of an engineered form of CD4
30 possessing a BamHI site just upstream of the membrane spanning domain (Aruffo et al., Proc. Natl. Acad. Sci. USA, 84:8573-8577 (1987b); Zettlmeissl et al., DNA Cell Biol., 9:347-353 (1990)) by standard techniques. To form

- 59 -

this fusion protein, a BamHI site may be engineered into the sequence at the appropriate location (again, by standard techniques). The gene fusions are introduced into a vaccinia virus expression plasmid (as described herein), and inserted into the genome of the vaccinia WR strain by homologous recombination and selection for growth in mycophenolic acid (Falkner et al., J. Virol., 62:1849-1854 (1988); Boyle et al., Gene, 65:123-128 (1988)). Flow cytometric analysis is used to examine expression by the vaccinia recombinants of the CD4-tyrosine kinase fusion proteins at the cell surface. Immunoprecipitation of cells infected with the vaccinia recombinants is used to confirm the results (as described above).

The efficacy of CD4 chimeras may be tested by any of the calcium mobilization or cytolysis assays described herein. In one particular example, a model target:effector system based on CD4 recognition of the HIV envelope gp120/gp41 complex is created. HeLa cells are infected with recombinant vaccinia viruses expressing gp120/gp41 (Chakrabarti et al., Nature, 320:535-537 (1986); Earl et al., J. Virol., 64:2448-2451 (1990)) and labeled with ⁵¹Cr. The labeled cells are incubated with cells from a human allospecific (CD8⁺, CD4⁻) cytotoxic T lymphocyte line which has been infected with vaccinia recombinants expressing the CD4-tyrosine kinase chimera, and examined to specific lysis.

To control for the possibility that vaccinia infection might promote artefactual recognition by CTL, similar cytolysis experiments are performed with target cells infected with vaccinia recombinants expressing the phosphatidylinositol linked form of CD16 (CD16_{PI}) and labeled with ⁵¹Cr, and with CTL infected with control recombinants expressing CD16 chimeras.

- 60 -

In another example, neutrophilic granulocytes, which have a very short lifespan ($\approx 4h$) in circulation and are intensely cytolytic, are attractive cells for expression of CD4-tyrosine kinase chimeras. Infection of
5 neutrophils with HIV is not likely to result in virus release, and the abundance of these cells (the most prevalent of the leukocytes) should facilitate host defense. Another attractive possibility for host cells are mature T cells, a population presently accessible to
10 retroviral engineering (Rosenberg, S.A. Sci. Am., 262:62-69 (1990)). With the aid of recombinant IL-2, T cell populations can be expanded in culture with relative ease, and the expanded populations typically have a limited lifespan when reinfused (Rosenberg et al., N.
15 Engl. J. Med., 323:570-578 (1990)).

Under the appropriate conditions, HIV recognition by cells expressing CD4 chimeras should also provide mitogenic stimuli, allowing the possibility that the armed cell population could respond dynamically to the
20 viral burden. Although we have focused here on the behavior of the fusion proteins in cytolytic T lymphocytes, expression of the chimeras in helper lymphocytes might provide an HIV-mobilized source of cytokines which could counteract the collapse of the
25 helper cell subset in AIDS. Recent description of several schemes for engineering resistance to infection at steps other than virus penetration (Friedman et al., Nature, 335:452-454 (1988); Green et al., Cell, 58:215-223 (1989); Malim et al., Cell, 58:205-214 (1989); Trono
30 et al., Cell, 59:113-120 (1989); Buonocore et al., Nature, 345:625-628 (1990)) suggests that cells bearing CD4 chimeras could be designed to thwart virus production by expression of appropriate agents having an intracellular site of action.

- 61 -

The ability to transmit signals to T lymphocytes through autonomous chimeras also provides the ability for the regulation of retrovirally engineered lymphocytes in vivo. Crosslinking stimuli, mediated for example by

5 specific IgM antibodies engineered to remove complement-binding domains, may allow such lymphocytes to increase in number in situ, while treatment with similar specific IgG antibodies (for example recognizing an amino acid variation engineered into the chimeric chain) could

10 selectively deplete the engineered population. We have previously determined that anti-CD4 IgM antibodies do not require additional crosslinking to mobilize calcium in Jurkat cells expressing CD4:ζ chimeras. The ability to regulate cell populations without recourse to repeated

15 extracorporeal amplification may substantially extend the range and efficacy of current uses proposed for genetically engineered T cells.

Other Embodiments

To create other chimeras consisting of protein

20 kinase intracellular sequences, cDNA or genomic sequences encoding an extracellular domain of the receptor can be endowed with a restriction site introduced at a location just preceding the extracellular domain of choice. The extracellular domain fragment terminating in the

25 restriction site can then be joined to the protein kinase sequences. Typical extracellular domains may be derived from receptors which recognize complement, carbohydrates, viral proteins, bacteria, protozoan or metazoan parasites, or proteins induced by them. Similarly,

30 ligands or receptors expressed by pathogens or tumor cells can be attached to protein kinase sequences to direct immune responses against cells bearing receptors recognizing those ligands.

To identify the minimal protein kinase sequences

35 necessary for cytolysis, a series of deletion mutants may

- 62 -

b prepared by standard techniques in which successively
more of the kinase intracellular domain is removed. Such
deletion mutants are tested for efficacy in any of the
assays described herein. Useful intracellular domains
5 for the Syk protein kinase include, for example, amino
acids 336-628 of the porcine Syk sequence and amino acids
338-630 of the human Syk sequence.

While the invention has been described in
connection with specific embodiments thereof, it will be
10 understood that it is capable of further modifications
and this application is intended to cover variations,
uses, or adaptations of the invention and including such
departures from the present disclosure as come within the
art to which the invention pertains and as may be applied
15 to the essential features hereinbefore set forth as
follows in the scope of the appended claims.

- 63 -

SEQUENCE LISTING**(1) GENERAL INFORMATION:**

(i) APPLICANT: Brian Seed
Charles Romeo
Waldemar Kolanus

(ii) TITLE OF INVENTION: REDIRECTION OF CELLULAR IMMUNITY
BY RECEPTOR CHIMERAS

(iii) NUMBER OF SEQUENCES: 8

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Fish & Richardson
(B) STREET: 225 Franklin Street
(C) CITY: Boston
(D) STATE: Massachusetts
(E) COUNTRY: U.S.A.
(F) ZIP: 02110-2804

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: 3.5" Diskette, 1.44 Mb
(B) COMPUTER: IBM PS/2 Model 50Z or 55SX
(C) OPERATING SYSTEM: MS-DOS (Version 5.0)
(D) SOFTWARE: WordPerfect (Version 5.1)

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: 08/093,210
(B) FILING DATE: July 16, 1993
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Paul T. Clark
(B) REGISTRATION NUMBER: 30,162
(C) REFERENCE/DOCKET NUMBER: 00786/195001

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (617) 542-5070
(B) TELEFAX: (617) 542-8906
(C) TELEX: 200154

- 64 -

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH:	33
(B) TYPE:	nucleic acid
(C) STRANDEDNESS:	single
(D) TOPOLOGY:	linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

CGCGGGGTGA CCGTGCCCTC CAGCAGCTTG GGC 33

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH:	50
(B) TYPE:	nucleic acid
(C) STRANDEDNESS:	single
(D) TOPOLOGY:	linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

CGCGGGGATC CGTCGTCCAG AGCCCGTCCA GCTCCCCGTC CTGGGCCTCA 50

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH:	33
(B) TYPE:	nucleic acid
(C) STRANDEDNESS:	single
(D) TOPOLOGY:	linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

CGCGGGCGGC CGCGACGCCG GCCAAGACAG CAC 33

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH:	33
(B) TYPE:	nucleic acid
(C) STRANDEDNESS:	single
(D) TOPOLOGY:	linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

CGCGTTGACG AGCAGCCAGT TGGGCAGCAG CAG 33

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 5:

- 65 -

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH:	630
(B) TYPE:	amino acid
(C) STRANDEDNESS:	N/A
(D) TOPOLOGY:	linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

Met	Ala	Asp	Ser	Ala	Asn	His	Leu	Pro	Phe	Phe	Phe	Gly	His	Ile	Thr
1				5					10					15	
Arg	Glu	Glu	Ala	Glu	Asp	Tyr	Leu	Val	Gln	Gly	Gly	Met	Ser	Asp	Gly
			20					25					30		
Leu	Tyr	Leu	Leu	Arg	Gln	Ser	Arg	Asn	Tyr	Leu	Gly	Gly	Phe	Ala	Leu
			35				40					45			
Ser	Val	Ala	His	Gly	Arg	Lys	Ala	His	Asn	Tyr	Thr	Ile	Glu	Arg	Glu
			50			55					60				
Leu	Asn	Gly	Thr	Tyr	Ala	Ile	Ala	Gly	Gly	Arg	Thr	His	Ala	Ser	Pro
65					70				75						80
Ala	Asp	Leu	Cys	Asn	Tyr	His	Ser	Gln	Glu	Ser	Asp	Gly	Leu	Val	Cys
			85						90					95	
Leu	Leu	Lys	Lys	Pro	Phe	Asn	Arg	Pro	Gln	Gly	Val	Gln	Pro	Lys	Thr
			100					105					110		
Gly	Pro	Phe	Glu	Asp	Leu	Lys	Glu	Asn	Leu	Ile	Arg	Glu	Tyr	Val	Lys
			115				120					125			
Gln	Thr	Met	Asn	Leu	Gln	Gly	Gln	Ala	Leu	Glu	Gln	Ala	Ile	Ile	Ser
			130			135					140				
Gln	Lys	Pro	Gln	Leu	Glu	Lys	Leu	Ile	Ala	Thr	Thr	Ala	His	Glu	Lys
145					150					155					160
Met	Pro	Trp	Phe	His	Gly	Lys	Ile	Ser	Arg	Glu	Ile	Ser	Thr	Gln	Ile
			165					170						175	
Val	Leu	Ile	Gly	Ser	Lys	Thr	Asn	Gly	Lys	Phe	Leu	Ile	Arg	Ala	Arg
			180					185					190		
Asp	Asn	Asn	Gly	Ser	Tyr	Ala	Leu	Cys	Leu	Leu	His	Ile	Gly	Lys	Val
			195				200					205			
Leu	His	Tyr	Arg	Ile	Asp	Lys	Asp	Lys	Thr	Gly	Lys	Leu	Ser	Ile	Pro
			210			215					220				
Glu	Gly	Lys	Lys	Phe	Asp	Thr	Leu	Trp	Gln	Leu	Val	Glu	His	Tyr	Ser
225					230					235					240
Tyr	Lys	Ala	Asp	Gly	Leu	Leu	Arg	Val	Leu	Thr	Val	Pro	Cys	Gln	Lys
			245						250					255	
Ile	Gly	Thr	Gln	Gly	Asn	Val	Asn	Phe	Gly	Gly	Arg	Pro	Gln	Leu	Phe
			260					265					270		
Gly	Ser	His	Pro	Ala	Thr	His	Ser	Ala	Gly	Gly	Ile	Ile	Ser	Arg	Ile
			275				280					285			
Lys	Ser	Tyr	Ser	Phe	Pro	Lys	Pro	Gly	His	Arg	Lys	Ser	Ser	Pro	Ala
			290			295					300				
Gln	Gly	Asn	Arg	Gln	Glu	Ser	Thr	Val	Ser	Phe	Asn	Phe	Tyr	Glu	Pro
305					310					315					320
Glu	Leu	Ala	Pro	His	Ala	Ala	Asp	Lys	Gly	Pro	Gln	Arg	Ile	Ala	Leu
			325						330					335	
Pro	Met	Asp	Thr	Glu	Val	Tyr	Glu	Ser	Pro	Tyr	Ala	Asp	Pro	Glu	Glu
			340					345					350		
Ile	Arg	Pro	Lys	Glu	Val	Tyr	Leu	Asp	Arg	Lys	Leu	Leu	Thr	Leu	Glu
			355				360					365			
Asp	Lys	Glu	Leu	Gly	Ser	Gly	Asn	Phe	Gly	Thr	Val	Lys	Lys	Gly	Tyr
			370			375					380				
Tyr	Gln	Met	Lys	Lys	Val	Val	Lys	Thr	Val	Ala	Val	Lys	Ile	Leu	Lys
385					390					395					400
Asn	Glu	Ala	Asn	Asp	Pro	Ala	Leu	Lys	Asp	Glu	Leu	Leu	Ala	Glu	Ala
			405						410					415	
Asn	Val	Met	Gln	Gln	Leu	Asp	Asn	Pro	Tyr	Ile	Val	Arg	Met	Ile	Gly
			420					425						430	

- 66 -

```

Ile Cys Glu Ala Glu Ser Trp Met Leu Val Met Glu Met Ala Glu Leu
      435      440      445
Gly Pro Leu Asn Lys Tyr Leu Gln Gln Asn Arg His Val Lys Leu Lys
      450      455      460
Asn Ile Ile Glu Leu Val His Gln Val Ser Met Gly Met Lys Tyr Leu
      465      470      475      480
Glu Glu Ser Asn Phe Val His Arg Asp Leu Ala Ala Arg Asn Val Leu
      485      490      495
Leu Val Thr Gln His Tyr Ala Lys Ile Ser Asp Phe Gly Leu Ser Lys
      500      505      510
Ala Leu Arg Ala Asp Glu Asn Tyr Tyr Lys Ala Gln Thr His Gly Lys
      515      520      525
Trp Pro Val Lys Trp Tyr Ala Pro Glu Cys Ile Asn Tyr Tyr Lys Phe
      530      535      540
Ser Ser Lys Ser Asp Val His Ser Phe Gly Val Leu Met Asn Glu Ala
      545      550      555      560
Phe Ser Tyr Gly Gln Lys Phe Tyr Arg Gly Met Lys Gly Ser Glu Val
      565      570      575
Ile Ala Met Leu Glu Lys Gly Glu Arg Met Gly Cys Pro Ala Gln Cys
      580      585      590
Pro Arg Glu Met Tyr Asp Leu Met Asn Leu Cys Trp Thr Tyr Asp Val
      595      600      605
Glu Asn Arg Pro Gly Phe Ala Ala Val Glu Leu Arg Leu Arg Asn Tyr
      610      615      620
Tyr Tyr Asp Val Val Asn
      625      630

```

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 6:

(i) SEQUENCE CHARACTERISTICS:

```

(A) LENGTH: 628
(B) TYPE: amino acid
(C) STRANDEDNESS: N/A
(D) TOPOLOGY: linear

```

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

```

Met Ala Asp Ser Ala Asn His Leu Pro Phe Phe Phe Gly Gln Ile Thr
  1      5      10      15
Arg Glu Glu Ala Glu Asp Tyr Leu Val Gln Gly Gly Met Ser Asp Gly
      20      25      30
Leu Tyr Leu Leu Arg Gln Ser Arg Asn Tyr Leu Gly Gly Phe Ala Leu
      35      40      45
Ser Val Ala Tyr Asp Arg Lys Ala His His Tyr Thr Ile Glu Arg Glu
      50      55      60
Leu Asn Gly Thr Tyr Ala Ile Ser Gly Gly Arg Thr His Gly Ser Pro
      65      70      75      80
Ala Glu Leu Cys His Tyr His Ser Gln Glu Leu Asp Gly Leu Val Cys
      85      90      95
Leu Leu Lys Asn Pro Phe Asn Arg Pro Pro Gly Val Gln Pro Lys Thr
      100      105      110
Gly Pro Phe Glu Asp Leu Lys Glu Asn Leu Ile Arg Glu Tyr Val Lys
      115      120      125
Gln Thr His Asn Leu Gln Gly Gln Ala Leu Glu Gln Ala Ile Ile Ser
      130      135      140
Gln Lys Pro Gln Leu Glu Lys Leu Ile Ala Thr Thr Ala His Glu Lys
      145      150      155      160
Met Pro Trp Phe His Gly Lys Ile Ser Arg Asp Glu Ser Glu Gln Ile
      165      170      175
Val Leu Ile Gly Ser Lys Ile Asn Gly Lys Phe Leu Ile Arg Ala Arg
      180      185      190

```

- 67 -

Asp Met Gly Ser Tyr Ala Leu Gly Leu Leu His Ile Gly Lys Val Leu
 195 200 205
 Met Tyr Arg Ile Asp Lys Asp Lys Thr Gly Lys Leu S r Ile Pro Gly
 210 215 220
 Gly Lys Asn Phe Asp Thr Leu Trp Gln Leu Val Glu Lys Tyr Ser Tyr
 225 230 235 240
 Lys Ser Asp Gly Leu Leu Arg Val Leu Thr Val Pro Cys Gln Lys Ile
 245 250 255
 Gly Gly Gln Thr Gly Asn Asp Ser Phe Arg Pro Gln Leu Phe Ser Ala
 260 265 270
 His Ser Ala Thr Trp Ser Ala Gly Ile Ile Ser Arg Ile Lys Ser
 275 280 285
 Tyr Ser Phe Pro Lys Pro Gly His Arg Lys Ala Ser Ser Pro Gln Gly
 290 295 300
 Asn Arg Pro Glu Ser Leu Val Ser Tyr Asn Phe Tyr Glu Ser Asp Arg
 305 310 315 320
 Gly Phe Trp Ala Asn Glu Arg Glu Ala Gln Arg Glu Ala Leu Pro Met
 325 330 335
 Asp Thr Glu Val Val Glu Ser Pro Tyr Ala Asp Pro Glu Glu Ile Arg
 340 345 350
 Pro Lys Glu Val Tyr Leu Asp Arg Lys Leu Leu Thr Leu Glu Asp Lys
 355 360 365
 Glu Leu Gly Ser Gly Asn Phe Gly Thr Val Lys Lys Gly Tyr Tyr Gln
 370 375 380
 Met Lys Lys Val Val Lys Thr Val Ala Val Lys Ile Leu Lys Asn Glu
 385 390 395 400
 Ala Asn Asp Pro Ala Leu Lys Asp Glu Leu Leu Ala Glu Ala Asn Val
 405 410 415
 Met Gln Gln Leu Asp Asn Pro Tyr Ile Val Arg Met Ile Gly Ile Cys
 420 425 430
 Glu Ala Ser Ser Trp Met Leu Val Met Glu Met Ala Glu Leu Gly Pro
 435 440 445
 Leu Asn Lys Tyr Leu Gln Gln Asn Arg His Val Lys Asp Lys Asn Ile
 450 455 460
 Ile Glu Leu Val His Gln Val Ser Met Gly Met Lys Tyr Leu Glu Glu
 465 470 475 480
 Cys Asn Pro Val His Arg Asp Leu Ala Ala Arg Asn Val Leu Leu Val
 485 490 495
 Thr Gln His Tyr Ala Lys Ile Ser Asp Phe Gly Leu Ser Lys Ala Leu
 500 505 510
 Arg Ala Asp Glu Asn Tyr Tyr Lys Ala Gln Thr His Gly Lys Trp Pro
 515 520 525
 Val Lys Trp Tyr Ala Pro Glu Cys Ile Asn Tyr Tyr Lys Phe Ser Ser
 530 535 540
 Lys Ser Asp Val Asn Ser Phe Gly Val Leu Met Trp Glu Ala Phe Ser
 545 550 555 560
 Tyr Gly Gln Lys Phe Tyr Arg Gly Met Lys Gly Ser Glu Val Ser Ala
 565 570 575
 Met Leu Glu Lys Gly Glu Arg Met Gly Cys Phe Phe Gly Cys Phe Arg
 580 585 590
 Glu Met Tyr Glu Leu Asn Thr Leu Cys Asn Thr Tyr Asp Val Glu Asn
 595 600 605
 Arg Pro Gly Phe Val Ala Val Glu Leu Arg Leu Arg Asn Tyr Tyr Tyr
 610 615 620

Asp Val Val Asn
 625

- 68 -

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH:	34
(B) TYPE:	nucleic acid
(C) STRANDEDNESS:	single
(D) TOPOLOGY:	linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

ATGGCAGACA GTGCCAACCA CTTGCCCTTC TTCT

34

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH:	39
(B) TYPE:	nucleic acid
(C) STRANDEDNESS:	single
(D) TOPOLOGY:	linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

CGCGGGGCGG CCGCTTTAAT TCACCACGTC GTAGTAGTA

39

- 69 -

Claims

1. A cell which expresses a membrane-bound, proteinaceous chimeric receptor, comprising (a) an intracellular portion of a protein-tyrosine kinase which
5 is capable of signalling said therapeutic cell to destroy a receptor-bound target cell or a receptor-bound target infective agent and (b) an extracellular portion which is capable of specifically recognizing and binding said target cell or said target infective agent, whereby each
10 of said therapeutic cells is capable of specifically recognizing and destroying said target cell or target infective agent.

2. The cell of claim 1, wherein said protein-tyrosine kinase is a member of the Syk kinase family.

15 3. The cell of claim 2, wherein said protein-tyrosine kinase is Syk.

4. The cell of claim 3, wherein said intracellular portion includes human Syk amino acids 336-628 or porcine Syk amino acids 338-630.

20 5. The cell of claim 1, wherein each of said therapeutic cells expresses a second membrane-bound, proteinaceous chimeric receptor, said second chimeric receptor comprising (a) an intracellular portion of a second protein-tyrosine kinase which is capable of
25 signalling said therapeutic cell to destroy a receptor-bound target cell or a receptor-bound target infective agent and (b) an extracellular portion which is capable of specifically recognizing and binding said target cell or said target infective agent, whereby each of said
30 therapeutic cells is capable of specifically recognizing and destroying a target cell or target infective agent.

- 70 -

6. The cell of claim 5, wherein one of said protein-tyrosine kinases is a member of the Syk kinase family and the other of said protein-tyrosine kinases is a member of the Src kinase family.

5 7. The cell of claim 6, wherein one of said protein-tyrosine kinases is ZAP-70 and the other of said protein-tyrosine kinases is Fyn.

8. The cell of claim 6, wherein one of said protein-tyrosine kinases is ZAP-70 and the other of said
10 protein-tyrosine kinases is Lck.

9. The cell of claims 7 or 8, wherein said ZAP-70 portion includes human ZAP-70 Tyr 369.

10. The cell of claims 1 or 5, wherein said therapeutic cells are selected from the group consisting
15 of: (a) T lymphocytes; (b) cytotoxic T lymphocytes; (c) natural killer cells; (d) neutrophils; (e) granulocytes; (f) macrophages; (g) mast cells; (h) HeLa cells; and (i) embryonic stem cells (ES).

11. The cell of claim 1, wherein said target
20 infective agent is an immunodeficiency virus.

12. The cell of claim 11, wherein said extracellular portion comprises an HIV envelope-binding portion of CD4.

13. The cell of claims 1 or 5, wherein said
25 binding is MHC-independent.

14. The cell of claims 1 or 5, wherein said extracellular portion comprises the ligand-binding

- 71 -

portion of a receptor, the receptor-binding portion of a ligand, the antigen-binding portion of an antibody, or a functional derivative thereof.

15. DNA encoding a chimeric receptor of claims 1
5 or 5.

16. A vector comprising the chimeric receptor DNA
of claims 1 or 5.

FIG. 1 A






FIG. 1 B



2/12

FIG. 1 C



3/12




FIG. 2

4/12




FIG. 3

5/12

FIG. 4 A





FIG. 4 B



6/12

FIG. 5 A




FIG. 5 B





FIG. 5 C



7/12

FIG. 6A





FIG. 6B



8/12

FIG. 7A

```

human 1  MADSAMSLPTTTPSTTAAAMSTLQSHHSDLTLLQSHSTLQSHVLLV 34
porcine 1  MADSAMSLPTTTPSTTAAAMSTLQSHHSDLTLLQSHSTLQSHVLLV 34

31  ANGLAAKNTTTEALSTTALAMSTAAAPALCTEDSODSLVCLLE 100
31  ATGAAKNTTTEALSTTALAMSTAAAPALCTEDSODSLVCLLE 100

101  PPTTPQGVQSTQSFEDLKLKIDSTVQSTVFLQCALQCALISQVPLE 130
101  PPTTPQGVQSTQSFEDLKLKIDSTVQSTVFLQCALQCALISQVPLE 130

131  ELIATTAALQSPWPHKISREISQIVLISSTVHGYLTAADHSTAL 200
131  ELIATTAALQSPWPHKISREISQIVLISSTVHGYLTAADHSTAL 200

201  CLLKQSVLSTEDIDSTVLSIPKHSSTLQSVLSTVSTHAKLLV 230
201  CLLKQSVLSTEDIDSTVLSIPKHSSTLQSVLSTVSTHAKLLV 230

231  TVPCQKISTQSGVVFQSPQLPQSHATWAGSISRLSTSTFVPH 297
231  TVPCQKISTQSGVVFQSPQLPQSHATWAGSISRLSTSTFVPH 297

300  KLSFACQKQSTVSTVSTPELAFWAGSPQKALPSTVSTVST 340
300  KLSFACQKQSTVSTVSTPELAFWAGSPQKALPSTVSTVST 340

340  KLSFACQKQSTVSTVSTPELAFWAGSPQKALPSTVSTVST 347
340  KLSFACQKQSTVSTVSTPELAFWAGSPQKALPSTVSTVST 347

350  PEXISPKVTLQKLLTLEKELQSHSTVSTVSTVSTVSTVSTV 380
350  PEXISPKVTLQKLLTLEKELQSHSTVSTVSTVSTVSTVSTV 387

400  EDLMDPALSCLLAAAPWQGLSWTYVWLSICLQSHSTVSTVST 440
400  EDLMDPALSCLLAAAPWQGLSWTYVWLSICLQSHSTVSTVST 447

450  PLKTYLQSHSTVSTVSTVSTVSTVSTVSTVSTVSTVSTVST 490
450  PLKTYLQSHSTVSTVSTVSTVSTVSTVSTVSTVSTVSTVST 497

500  QNTARISQFSLKALADSTTACTHSDVSTVSTVSTVSTVST 540
500  QNTARISQFSLKALADSTTACTHSDVSTVSTVSTVSTVST 547

550  VSTVSTVSTVSTVSTVSTVSTVSTVSTVSTVSTVSTVST 590
550  VSTVSTVSTVSTVSTVSTVSTVSTVSTVSTVSTVSTVST 597

600  NGLCTTSTVSTVSTVSTVSTVSTVSTVSTVSTVSTVST 630
600  NGLCTTSTVSTVSTVSTVSTVSTVSTVSTVSTVSTVST 637

650  NGLCTTSTVSTVSTVSTVSTVSTVSTVSTVSTVSTVST 680
650  NGLCTTSTVSTVSTVSTVSTVSTVSTVSTVSTVSTVST 687

```

9/12

FIG. 7 B





FIG. 7 C



10/12




FIG. 8

11/12



FIG. 9


12/12

FIG. 10 A



CD16/7/syk

FIG. 10 B



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/06675

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : C12N 5/10, 15/00, 15/11

US CL : 435/240.1, 240.2, 320.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/240.1, 240.2, 320.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS; DIALOG DATABASES: MEDLINE, BIOSIS, CA SEARCH, WORLD PATENT INDEX, AIDSLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO, A, 92/15322 (SEED ET AL.) 17 September 1992, see entire document.	1-16
Y	Cell, Volume 64, issued 08 March 1991, Romeo et al., "Cellular Immunity to HIV Activated by CD4 Fused to T Cell or Fc Receptor Polypeptides," pages 1037-1046, see entire document.	1-16
Y	Cell, Volume 71, issued 13 November 1992, Chan et al., "ZAP-70: a 70 kd Protein-Tyrosine Kinase That Associates with the TCR ζ Chain," pages 649-662, see entire document.	1-16

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

31 AUGUST 1994

Date of mailing of the international search report

14 OCT 1994

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JOHNNY F. RAILEY II, PH.D.

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/06675

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Cold Spring Harbor Symposia on Quantitative Biology, Volume 57, issued 1992, Weiss et al., "Regulation of Protein Tyrosine Kinase Activation by the T-cell Antigen Receptor ζ Chain," pages 107-116, see entire document.	1-16
Y	Cold Spring Harbor Symposia on Quantitative Biology, Volume 57, issued 1992, Romeo et al., "Activation of Immune System Effector Function by T-cell or Fc Receptor Intracellular Domains," pages 117-125, see entire document.	1-16
Y	Molecular and Cellular Biology, Volume 12, Number 12, issued December 1992, Timson Gauen et al., "p59fyn Tyrosine Kinase Associates with Multiple T-Cell Receptor Subunits through Its Unique Amino-Terminal Domain," pages 5438-5446, see entire document.	1-16